

A small variation of the Taylor Method and periodic solutions of the 3-body problem

Oscar Perdomo

Abstract

In this paper we define a small variation of the Taylor method and a formula for the global error of this new numerical method that allows us to keep track of the round-off error and does not require previous knowledge of the exact solution. As an application we provide a rigorous proof of the construction/existence of a periodic solution of the three body problem. Some images of this periodic motion can be seen at <https://www.youtube.com/watch?v=fSmQyeKcj5k>




Figure 0.1: This is the orbit of one of the bodies in the periodic solution that we have mathematically shown to exist. The three masses are 100, 100 and 200 and the period of this solution is $T = 96.241 \dots$. We are assuming that the gravitational constant is 1.

1 Introduction

1.1 The numerical Part

When we think about using a numerical method to do a mathematical proof, the first thing that comes to our mind is that we need to consider the error of the numerical method. Very soon we realize that the standard formula for the error is no very useful due to the fact that it assumes that all the basic operations are being made with no error and this is computationally very expensive. As an example, if we consider the initial value problem $y' = y - \frac{1}{3}y^2$ with $y(0) = \frac{1}{2}$ and we want to estimate $y(\frac{3}{10})$ using the Euler Method with $h = \frac{1}{100}$, we see that, even though we are considering the computational cheapest numerical method and we are only doing 30 iterations, a regular Computer Algebra System (Mathematica 10 in this case) will need 8696.99 seconds to do these 30 iterations most likely because the final answer is a rational number of the form $\frac{p}{q}$ with p and q integers both with 2607760525 digits, more than 2.6 billions digits. On the other hand, if we allow the Computer Algebra System to have round-off error in every operation involved in each iteration, it becomes challenging to keep track of the error because easily, each iteration may have a few dozens of operations. In our example above, each iteration has 3 operations: one raising to the square, one product and a difference. We will exploit the fact that most of the Computer Algebra Systems (CAS) can compute, with a mathematical precision, the two integers i and j such that $i \leq r_1 + \sqrt{r_2} \leq j$ where r_1 and r_2 rational number and $r_2 \geq 0$. The new numerical method that we are proposing in this paper allows us to work all the time with mathematical precision without paying the price of dealing with numbers that have huge expression using integers. Our numerical method will do the operations in each iteration with a mathematical precision and then at the end of the iteration, it will find with a mathematical precision a rational number that approximates the output of the iteration within a distance H . In order to be able to use the method, we find a formula for the error between the real value of the solution of the ODE and the approximation given by the numerical method in terms of the two values: h , the desired value for the step, and H , the desired value for the rounding in each iteration. This is done in Theorem (2.3). In general, when we try to estimate the error of a numerical method, a problem that we face is that we need to have some a-priori bounds of the solution of the differential equation that we do not know. An important aspect of Theorem (2.3) is that it does not need previous knowledge of the exact solution.

1.2 Periodic solutions of the three body problem

Poincare showed that the three body problem has a chaotic behavior that makes it difficult to solve. For this reason, it is not surprising that the only explicit solutions were discovered more than 240 years ago by Euler in 1765 and by Lagrange in 1772. Recall that we have a periodic solution when the values of the positions and the velocities of the three bodies, after some time T , agree with the values of the positions and velocities at $t = 0$. Usually, when the values of a solution after T are within a small distance ϵ from the initial condition, this solution is called *numerically periodic*. There is an enormous amount of numerically periodic solutions, for example, in 1975 [5], H. Henon showed a family of numerically periodic planar solutions of the three body with $\epsilon = 10^{-14}$. Despite the abundance of numerically periodic solutions, the task of showing that numerically periodic solutions are periodic is a difficult one. In 2000, Chenciner and Montgomery [1] showed that a numerically solution found by Moore in 1993, [4], was indeed periodic. This example represents (to my knowledge) the first example of a periodic solution that has a numerical image associated with it and does not have an explicit formula. It is important to mention that proofs showing the existence of periodic solutions have been found before, for example, Meyer and Schmidt1993 [3]. These solutions are usually near a bifurcation point of a family of solutions described with explicit formulas. After the paper by Chenciner and Montgomery, there has been more proofs showing that some numerical solutions are periodic. Among them, we have papers by Terracini and Ferrario, K.C Chen and Simo and Kapitza and Gronchi. An excellent account of the work done in this direction so far can be found in the site <http://montgomery.math.ucsc.edu/Nbdy.html>

In this paper we will show that the numerical periodic solution given by the initial condition explained in Figure 1.1 is periodic.

To describe the motion, let us assume that t represents seconds. Keep in mind that the units of time, mass and distance, have been adjusted so that the gravitational constant is 1. For this periodic solution the motion starts with two bodies, each one with a mass of 100, separated 20 units and the third body with a mass of 200 right in the middle. If we reference the body in the middle as body 1, then, we have that the initial position of the body 1 is $(0, 0, 0)$ and the initial position of the other two bodies are $(10, 0, 0)$ and $(-10, 0, 0)$. Body 1 will always stay in the z -axis while the other two bodies will move around the z -axis. During the first $t_0 = 2.673 \dots$ seconds, Body 1 moves straight up to the position $(0, 0, 2.453 \dots) = (0, 0, z_1)$, this is the farthest up this body will get. Simultaneously, the other two bodies move down and around the z -axis; after these first t_0 seconds they are both z_1 units below the x - y plane, they are $r_1 = 9.43 \dots$ units away from the the z axis; and they have made




Figure 1.1: There is a periodic solution with period $T = 96.241\dots$, $a = 4.3170\dots$ and $b = 1.4903\dots$.

a $\frac{7\pi}{18}$ -rotation (70 degrees) with respect to the z -axis. At every instance, the distance to the z -axis of these two bodies is the same and it is twice the distance between them. In this way, after the first t_0 seconds Body 2 is at $(r_1 \cos \frac{7\pi}{18}, r_1 \sin \frac{7\pi}{18}, -z_1)$ and Body 3 is at $(-r_1 \cos \frac{7\pi}{18}, -r_1 \sin \frac{7\pi}{18}, -z_1)$. During the next t_0 seconds the body 1 moves down getting back to the origin and the other two bodies will simultaneously rotate, move up and move apart from the z axis; they will rotate another $\frac{7\pi}{18}$, and they will go back to be 10 units apart from the z -axis. Notice that after $2t_0$ seconds the relative positions with respect to each other are the same as in the starting position. The only difference is that Body 1 is now going down and the other two bodies are going up. With respect to a fixed reference frame, after $2t_0$ seconds, the positions of all three bodies differ from the starting position by a rotation of $\frac{7\pi}{9}$. The next t_0 seconds Body 1 will reach its lowest point $(0, 0, -z_1)$ while the other two bodies will reach the highest point rotating another $\frac{7\pi}{18}$. Finally after another t_0 seconds the relative positions with respect to each other are the same as the starting position, and this time the body 1 is going up as it was at the starting position. With respect to a fixed frame, the positions after $4t_0$ seconds differ from the starting position by a rotation of $\frac{14\pi}{9}$. Doing 9 more of these $4t_0$ cycles will bring the three bodies to the starting position. In this process, bodies 2 and 3 will have completed 7 rotations around the z -axis. The trajectory of Body 2 is shown in Figure 0.1.

Each color represents a cycle of $4t_0$ seconds, for this reason there are 9 colors in the picture. Body 3 does not share the same trajectory as Body 2. The trajectory of Body 3 is the reflection with respect to the x - y plane of the trajectory of the body 2.

We would like to emphasize that the goal of this part of the paper is not to show the existence of this type of solutions: This was done by Meyer and Schmidt [3], nor to show numerically periodic solutions of this type: We can see similar images of solutions like the one we are showing (when the three masses are the same), in the work by Yan and Ouyang [7]. The goal of this part of the paper is to give a rigorous proof that a particular numerically periodic solution is indeed periodic.

We will reduce the proof of the periodicity of this solution to show that three functions defined in an open set of R^3 must vanish simultaneously. The variables in the domain of these three functions are given by triples (t, a, b) where a and b are explained in Figure 1.1

The author has found some numerical solutions of the same type, not only for the 3 body problem but for the n -body problem. Some aspects of these solutions have been posted online:

The link <https://www.youtube.com/watch?v=PtEMb6Rvflg> shows a periodic solution of the 6 body problem.

The link <https://www.youtube.com/watch?v=2Wpv6vpOxXk> shows a periodic solution of the four body problem.

The link <https://www.youtube.com/watch?v=hjQp1P09560> shows a periodic solution of the three body problem.

The images in the videos were generated by solving the differential equation that governs the n body problem and then posting 10 pictures per second, in this way, the time in the video is proportional to the time t of the solution.

The technique used in this paper to prove periodicity is different to the one used by Chenciner and Montgomery, they used variational methods. There are five ingredients in our proof that the solution that we are considering is periodic: (i) The Poincare-Miranda theorem, Theorem 4.4, which is essentially a generalization of the intermediate value theorem. (ii) A symmetry result that allow to integrate the ordinary differential equation (ODE) over a quarter of a period instead of the whole period. Lemma 6.2. (iii) The Round Taylor Method to solve differential equations, that allow us to estimate the values of the functions that we are considering. Section 2. (iv) A lemma related with the the implicit function theorem that allow us to find a set where the solution of a system of equations of the form

$f_1(t, a, b) = 0, f_2(t, a, b) = 0$ is given by only points on a connected smooth curve, Section 3. (v) A theorem that tell us how to compute the partial derivative of an ODE with respect to the initial conditions and parameters in the ODE. [2].

The author would like to express his gratitude to Andrés Rivera, Richard Montgomery, Carles Simo and Marian Anton for their valuable comments on this work.

2 The Round Taylor Method

2.1 The Round Taylor Method

We will be using a small modification of the Taylor method of order m to estimate solutions of differential equations. This modification will allow the computer that implements the method to work all the time with rational number and its square roots instead of working with approximations of numbers. In this way, the method is free of error coming from decimal approximation, this is, this method is *round -off* error free. Let us call this method the *Round Taylor Method*. Assume we have the ordinary differential equation

$$Y'(t) = f(Y(t)), \quad Y(0) = y_0 \quad (2.1)$$

where $Y(t) = (Y_1(t), \dots, Y_n(t))^T$ and $f(y) = (f_1(y), \dots, f_n(y))^T$. Here v^T stands for the transpose of the vector v . Given two positive rational number h and H , we define the sequence of points in R^n that starts with $z_0 = y_0$ and follows using the recursive formulas

$$\begin{aligned} y_1 &= z_0 + f(z_0)h + F_1(z_0)\frac{h^2}{2!} + \dots F_{m-1}(z_0)\frac{h^m}{m!} \\ z_1 &= \text{a rational number such that } |z_1 - y_1| \leq H \\ y_2 &= z_1 + f(z_1)h + F_1(z_1)\frac{h^2}{2!} + \dots F_{m-1}(z_1)\frac{h^m}{m!} \\ &\vdots \\ z_i &= \text{a rational number such that } |z_i - y_i| \leq H \\ y_{i+1} &= z_i + f(z_i)h + F_1(z_i)\frac{h^2}{2!} + \dots F_{m-1}(z_i)\frac{h^m}{m!} \end{aligned}$$

where $F_1 = Dff$, $F_2 = DF_1f$, $\dots$ $F_{m-1} = DF_{m-2}f$. Here D denotes the derivative operator that takes a function from R^n to R^n to the matrix which columns are the

partial derivatives with respect to the variables. In order to define z_i from y_i , we may use -and will use in this paper- the Floor function $\lfloor x \rfloor$ that assigns to a real number x the largest integer not greater than x . When $x = (x_1, \dots, x_k)$ is a vector, $\lfloor x \rfloor = (\lfloor x_1 \rfloor, \dots, \lfloor x_k \rfloor)$. It is clear that a possible choice for z_i is $z_i = H \lfloor \frac{1}{H} y_i \rfloor$.

As an example, when we use the Round Taylor method of order 1 for the differential equation given in the introduction, this is, $n = 1$, $f(y) = y - \frac{y^2}{3}$, $y_0 = \frac{1}{2}$. When $h = 1/100$ and $H = 10^{-6}$, we obtain that $y_0 = 1/2 = z_0$ and the next 10 values of $\{y_i, z_i\}$ are given by

$$\begin{pmatrix} \frac{121}{240} & \frac{252083}{500000} \\ \frac{38127028661111}{75000000000000} & \frac{12709}{25000} \\ \frac{96109156319}{187500000000} & \frac{256291}{500000} \\ \frac{38762401423319}{75000000000000} & \frac{16151}{31250} \\ \frac{152668926449}{292968750000} & \frac{521109}{1000000} \\ \frac{52541490803373}{100000000000000} & \frac{262707}{500000} \\ \frac{13243698510717}{25000000000000} & \frac{529747}{1000000} \\ \frac{160232709115991}{300000000000000} & \frac{534109}{1000000} \\ \frac{161549754576119}{300000000000000} & \frac{538499}{1000000} \\ \frac{162875215826999}{300000000000000} & \frac{542917}{1000000} \end{pmatrix}$$

Let us start the process of finding the formula for the error of this numerical method.

Lemma 2.1. *The sequence given by the recursive formula $q_{i+1} = (1+L)q_i + p$ (with $L \neq 0$) satisfies that $q_k = \frac{p+Lq_0}{L}(1+L)^k - \frac{p}{L}$.*

Proof. The proof follows by induction. Clearly the formula works for $k = 0$. Now, assuming the formula works for $k \geq 0$, we have:

$$q_{k+1} = (1+L)q_k + p = (1+L) \left(\frac{p+Lq_0}{L}(1+L)^k - \frac{p}{L} \right) + p = \frac{p+Lq_0}{L}(1+L)^{k+1} - \frac{p}{L}$$

This finishes the proof of the lemma. $\square$

There are different norms that we can use in the set of matrices, in order to establish the the one that we are using, we state the following lemma.

Lemma 2.2. *Let A be an $n \times n$ matrix, if $|A|^2$ denote the sum of the square of its entries, then for any vector v in R^n we have that $|Av| \leq |A||v|$.*

Proof. If A^i denote the columns of A and v_i denote the entries of v , then we have that

$$|Av| = \left| \sum_{i=1}^n v_i A^i \right| \leq \sum_{i=1}^n |v_i| |A^i| \leq |A||v|$$

$\square$

Theorem 2.3. *Let h and H be two positive numbers and k a positive integer. Let us consider the sequences $\{(y_i, z_i)\}_{i=0}^k$ given by the Round Taylor Method associated with the values h and H for the ODE $Y'(t) = f(Y(t))$, $Y(0) = y_0$, described in the beginning of this section. Let us assume that we can find constants $M_0, M_1, \dots, M_n, K_0, \dots, K_{m-1}$ and sets $U_1 = \{(u_1, \dots, u_n) : b_1 \leq u_1 \leq c_1, \dots, b_n \leq u_n \leq c_n\}$ and $U_2 = \{(u_1, \dots, u_n) : b_1 - \epsilon < u_1 < c_1 + \epsilon, \dots, b_n - \epsilon < u_n < c_n + \epsilon\}$ with $\epsilon > M_0 h + \tilde{H}$, where $\tilde{H} = \frac{M \frac{h^m}{(m+1)!} + \frac{H}{h}}{L} (e^{Lkh} - 1)$ and $M = \sqrt{M_1^2 + \dots + M_n^2}$ and $L = K_0 + K_1 \frac{h}{2!} + \dots + K_{m-1} \frac{h^{m-1}}{m!}$. If*

- *The map f and all its partial derivative with order less than $m+2$ are continuous in an open set that contain the closure of U_2 .*
- *$z_j \in U_1$ for $j = 0, \dots, k$.*
- *$|F_{mi}(u)| \leq M_i$ for $i = 1, \dots, n$, for all $u \in U_2$. Here $F_m = (F_{m1}, \dots, F_{mn})$.*
- *$|Df(u)| \leq K_0$ and $|DF_i(u)| \leq K_i$ for $i = 1, \dots, m-1$ and $u \in U_2$.*
- *$|f_i(u)| \leq M_0$ for $i = 1, \dots, n$ and $u \in U_2$.*

then, the solution $Y(t)$ of the system of ordinary differential equation is defined on $[0, hk]$ and for any positive integer $j \leq k$, we have that $|Y(jh) - z_j| \leq \tilde{H}$.

Proof. Let us start by checking that for any pair of points u_1 and u_2 in U_2 we have that $|f(u_2) - f(u_1)| \leq K_0|u_2 - u_1|$ and $|F_i(u_2) - F_i(u_1)| \leq K_i|u_2 - u_1|$ for $i = 1, \dots, m-1$. We have that

$$\begin{aligned} |f(u_2) - f(u_1)| &= \left| \int_0^1 \frac{df(u_1 + t(u_2 - u_1))}{dt} dt \right| \\ &= \left| \int_0^1 Df(u_1 + t(u_2 - u_1)) (u_2 - u_1) dt \right| \leq K_0|u_2 - u_1| \end{aligned}$$

The proof of the inequalities $|F_i(u_2) - F_i(u_1)| \leq K_i|u_2 - u_1|$, for $i = 1, \dots, m-1$ is similar. For any non negative integer $l \leq k$ such that lh is in the domain of Y let us denote by $x_l = Y(lh)$. Notice that anytime $Y(t)$ is in the closure of U_2 for all $t \in [0, t_1]$, then it is possible to extend $Y(t)$ to an interval of the form $[0, t_1 + \delta]$ with $\delta > 0$. We will show that for any $l \leq k$, $Y(t)$ is defined on $[0, lh]$, $Y(t) \in U_2$ for all $t \in [0, hl]$ and $|Y(lh) - z_l| \leq \tilde{H}$. Clearly the result hold for $l = 0$ because $|x_0 - z_0| = 0$. Let us assume that $Y(t)$ is defined for all $t \in [0, jh]$ with $Y(t) \in U_2$ and $|x_l - z_l| < \tilde{H}$ for all $0 \leq l \leq j < k$, we will prove the theorem by showing that $Y(t)$ is defined for all $t \in [0, (j+1)h]$ with $Y(t) \in U_2$ and $|x_{j+1} - z_{j+1}| < \tilde{H}$. Let us show that for any $t \in [jh, (j+1)h]$, $Y(t) \in U_2$. Since $Y(t)$ is continuous and U_2 is open, then $Y(hj + \tau) \in U_2$ for small positive values of τ . Let us show by contraction that $Y(hj + \tau) \in U_2$ for all $\tau \in [0, h]$. If $Y(hj + \tau) \notin U_2$ for all $\tau \in [0, h]$, then we can find τ^* such that $Y(jh + \tau) \in U_2$ for all $\tau \in (0, \tau^*)$ and $Y(jh + \tau^*) \notin U_2$. Therefore, writing $Y = (Y_1, \dots, Y_n)$, we can find a positive integer $w \leq n$ such that either $Y_w(jh + \tau^*) = b_w - \epsilon$ or $Y_w(jh + \tau^*) = c_w + \epsilon$ and $Y_e(jh + \tau) \in (b_e - \epsilon, c_e + \epsilon)$ for all positive integers $e \leq n$ and all $\tau \in (0, \tau^*)$. Denoting $z_j = (z_{j_1}, \dots, z_{j_n})$, we have that for some $\xi \in (0, \tau^*)$,

$$\begin{aligned} |Y_w(jh + \tau^*) - z_{j_w}| &\leq |Y_w(jh + \tau^*) - Y_w(jh)| + |Y_w(jh) - z_{j_w}| \\ &\leq |\dot{Y}_w(hj + \xi)|\tau^* + \tilde{H} \\ &\leq |f_w(Y(jh + \xi))|\tau^* + \tilde{H} \\ &< M_0h + \tilde{H} < \epsilon \end{aligned}$$

Since $z_j \in U_1$, the inequality above contradicts the fact that either $Y_w(jh + \tau^*) = b_w - \epsilon$ or $Y_w(jh + \tau^*) = c_w + \epsilon$. This contradiction shows that $Y(t) \in U_2$ for all $t \in [0, (j+1)h]$, in particular $x_{j+1} \in U_2$. Let us prove now that $|x_{j+1} - z_{j+1}| < \tilde{H}$.

For any $i = 0, \dots, j+1$ we have that

$$\begin{aligned}
|x_{i+1} - z_{i+1}| &\leq |x_{i+1} - y_{i+1}| + |y_{i+1} - z_{i+1}| \\
&\leq |x_i + f(x_i)h + F_1(x_i)\frac{h^2}{2!} + \dots + F_{m-1}(x_i)\frac{h^m}{m!} - y_{i+1}| + \frac{Mh^{m+1}}{(m+1)!} + H \\
&\leq |x_i - z_i|(1 + hL) + \frac{Mh^{m+1}}{(m+1)!} + H
\end{aligned}$$

Let us define $p = \frac{Mh^{m+1}}{(m+1)!} + H$, $q_0 = 0$ and $q_{i+1} = (1 + hL)q_i + p$. By induction we can show that $|x_i - z_i| \leq q_i$ for all $i \leq j+1$. Using Lemma (2.1) we obtain that

$$\begin{aligned}
|Y((j+1)h) - z_{j+1}| &= |x_{j+1} - z_{j+1}| \leq \frac{p}{hL} ((1 + hL)^{j+1} - 1) \\
&\leq \frac{p}{hL} (e^{Lhk} - 1) = \tilde{H}
\end{aligned}$$

This finishes the proof. □

The following theorem is well known. A reference for the particular case when f and g are Lipschitz real value functions can be found at Earl A. Coddington, An Introduction to ordinary differential equations -Dover Publication - 1989. A reference for a more general case can be found at Herbert Amann, Ordinary Differential Equations An Introduction to Nonlinear Analysis - Walter de Gruyter - 1990.

Theorem 2.4. *Let us assume that $f, g : U \subset \mathbb{R}^n \rightarrow \mathbb{R}^n$ are C^1 functions defined on an open convex set U such that $|Df(u)| < K$, $|Dg(u)| < K$ and $|f(u) - g(u)| < \epsilon$. If $y(t)$ and $z(t)$ satisfy $\dot{y}(t) = f(y(t))$ and $\dot{z}(t) = g(z(t))$ respectively, then*

$$|y(t) - z(t)| \leq |y(t_0) - z(t_0)|e^{K|t-t_0|} + \frac{\epsilon}{K} (e^{K|t-t_0|} - 1)$$

3 On the implicit function theorem

Let us consider the set $\Sigma = \{x = (x_1, x_2, x_3) : f_1(x) = 0, f_2(x) = 0\}$ where f_1 and f_2 are real value functions defined on an open set of $\mathbb{R}^3$. By the Implicit Function Theorem we know that if $p_0 \in \Sigma$ and $\nabla f_1(p_0) \times \nabla f_2(p_0)$ is not the zero vector then there exists an open set U that contains p_0 such that $\Sigma \cap U$ is given a

regular connected curve. The following lemma give us an estimate on how big this open set U can be, under the assumption that we know that there is point in Σ in a small box with dimensions $2\mu_1$, $2\mu_2$ and $2\mu_3$.

Theorem 3.1. *Let us assume that $f_1, f_2 : R^3 \rightarrow R$ are smooth functions, $\mu_1, \mu_2, \mu_3, \delta_1, \epsilon_1, \epsilon_2, \tilde{\epsilon}_1, \tilde{\epsilon}_2, \delta_2, \tilde{\delta}_1, \tilde{\delta}_2$ are positive numbers such that for $i = 1, 2$, $\mu_i < \epsilon_i < \delta_i$, $\rho_i = \frac{\tilde{\epsilon}_i - \mu_i}{m_i} - \mu_3 > 0$ where $m_1 = \frac{\epsilon_1(\tilde{\delta}_2 + \epsilon_2)}{\delta_1\delta_2 - \epsilon_1\epsilon_2}$ and $m_2 = \frac{\epsilon_2(\tilde{\delta}_1 + \epsilon_1)}{\delta_1\delta_2 - \epsilon_1\epsilon_2}$ and*

$$U = \{(x, y, z) : |x| < \tilde{\epsilon}_1, |y| < \tilde{\epsilon}_2, |z| < \rho\} \text{ where } \rho < \min\{\rho_1, \rho_2\}.$$

If there exists $p_0 = (x_0, y_0, z_0)$ such that $f_1(p_0) = f_2(p_0) = 0$ with $|x_0| < \mu_1$, $|y_0| < \mu_2$, $|z_0| < \mu_3$ and for every $p \in U$ we have that

$$\delta_1 < \left| \frac{\partial f_1}{\partial x_1} \right| < \tilde{\delta}_1, \left| \frac{\partial f_1}{\partial x_2} \right| < \epsilon_1, \left| \frac{\partial f_1}{\partial x_3} \right| < \epsilon_1, \delta_2 < \left| \frac{\partial f_2}{\partial x_2} \right| < \tilde{\delta}_2, \left| \frac{\partial f_2}{\partial x_1} \right| < \epsilon_2, \left| \frac{\partial f_2}{\partial x_3} \right| < \epsilon_2$$

then, for every z_1 with $|z_1| < \rho$, there exists a unique (x_1, y_1) such that $(x_1, y_1, z_1) \in U$ is a solution of the equations $f_1(x, y, z) = 0$, $f_2(x, y, z) = 0$.

Proof. Let us denote by $\xi = (\xi_1, \xi_2, \xi_3) = \nabla f_1 \times \nabla f_2$. Direct computations show that $|\xi_1| \leq \epsilon_1(\tilde{\delta}_2 + \epsilon_2)$, $|\xi_2| \leq \epsilon_2(\tilde{\delta}_1 + \epsilon_1)$, and $|\xi_3| \geq \delta_1\delta_2 - \epsilon_1\epsilon_2$. Let $\alpha(t) = (x(t), y(t), z(t))$ be the integral curve of the vector field ξ such that $\alpha(0) = p_0$. Since $|\xi_3| > 0$ we have that the vector field ξ never vanishes on U , therefore there exist $T_2 > 0$ and $T_1 < 0$ such that $\alpha(t) \in U$ for all $t \in (T_1, T_2)$ and $\alpha(T_1)$ and $\alpha(T_2)$ are in the boundary of U , in particular we have that either $x(T_2) = \pm\tilde{\epsilon}_1$, $y(T_2) = \pm\tilde{\epsilon}_2$ or $z(T_2) = \pm\rho$. We will prove that $z(T_2)$ must be either ρ or $-\rho$. Using the fact that $\dot{x} = \xi_1(\alpha(t))$, $\dot{y} = \xi_2(\alpha(t))$ and $\dot{z} = \xi_3(\alpha(t))$ we obtain that,

$$\left| \frac{\dot{x}}{\dot{z}} \right| \leq \frac{\epsilon_1(\tilde{\delta}_2 + \epsilon_2)}{\delta_1\delta_2 - \epsilon_1\epsilon_2} = m_1 \quad \text{and} \quad \left| \frac{\dot{y}}{\dot{z}} \right| \leq \frac{\epsilon_2(\tilde{\delta}_1 + \epsilon_1)}{\delta_1\delta_2 - \epsilon_1\epsilon_2} = m_2 \quad (3.1)$$

Recall that $z(t)$ is one to one because its derivative never vanishes. Let us denote by h the inverse function of z . We have that $h(z(t)) = t$. If $g_1(\tau) = x(h(\tau))$, then $g_1(z_0) = x_0$ and $|g_1'(\tau)| < \frac{\epsilon_1(\tilde{\delta}_2 + \epsilon_2)}{\delta_1\delta_2 - \epsilon_1\epsilon_2}$. Therefore,

$$|x(T_2) - x_0| = |g(z(T_2)) - g_1(z_0)| = |g'(\tau^*)||z(T_2) - z_0| \leq m_1|z(T_2) - z_0| + m_1\mu_3$$

and then $|x(T_2)| < \mu_1 + m_1\rho_1 + m_1\mu_3 = \tilde{\epsilon}_1$. Likewise we can show that $|y(T_2)| < \tilde{\epsilon}_2$. Therefore we must have $|z(T_2)| = \rho$. The same arguments show that $|z(T_1)|$ must be either ρ or $-\rho$. Since the function $\dot{z}(t) \neq 0$ and $[T_1, T_2]$ is connected then we get that for every z_1 with $|z_1| < \rho$, there exists a $p_1 = (x_1, y_1, z_1) = \alpha(t_1) \in U$ such that $f_1(p_1) = 0$ and $f_2(p_1) = 0$. Let us prove that p_1 is unique. If $p_2 = (x_2, y_2, z_1) \in U$ and $p_2 \neq p_1$, then either $|x_2 - x_1| \leq |y_2 - y_1| \neq 0$ or $|y_2 - y_1| \leq |x_2 - x_1| \neq 0$. If $|x_2 - x_1| \leq |y_2 - y_1| \neq 0$. Since we are assuming that $f_2(x_1, x_2, x_3) = 0$ then,

$$\begin{aligned} |f_2(x_2, y_2, z_1)| &\geq |f_2(x_2, y_2, z_1) - f_2(x_2, y_1, z_1)| - |f_2(x_2, y_1, z_1) - f_2(x_1, y_1, z_1)| \\ &\geq |y_2 - y_1|\delta_2 - |x_2 - x_1|\epsilon_2 > 0 \end{aligned}$$

Likewise we can show that if $|y_2 - y_1| \leq |x_2 - x_1| \neq 0$, then $|f_1(p_2)| > 0$. Therefore the only solution of the system $f_1(p) = f_2(p) = 0$ on the set U are those points in the curve $\alpha : [T_1, T_2] \rightarrow U$. This finishes the proof. $\square$

For any $u = (u_1, u_2, u_3)$ and $v = (v_1, v_2, v_3)$, let us denote by $u \cdot v = u_1v_1 + u_2v_2 + u_3v_3$ the dot product of u and v . Given a function $f : R^3 \rightarrow R$, we denote $D_v f_i = \nabla f_i \cdot v$. By doing an orthogonal change of coordinates we have the following corollary,

4 The differential equation, a symmetry result and the Poincare Miranda Theorem

Let us start introducing the differential equation for the subfamily of solutions of the three body that we are considering in this paper.

Proposition 4.1. *If f , r and θ satisfy the initial value system of differential equations*

$$\ddot{f} = -\frac{400}{s^3}f, \quad \ddot{r} = \frac{100a^2}{r^3} - \frac{25}{r^2} - \frac{200r}{s^3}, \quad r^2\dot{\theta} = 10a, \quad (4.1)$$

with $r(0) = 10$, $f(0) = 0$, $\dot{r}(0) = 0$, $\dot{f}(0) = b$, $\theta(0) = 0$, where $s = \sqrt{r^2 + 4f^2}$ then,

$$\begin{aligned}
x(t) &= (0, 0, f(t)) \\
y(t) &= (r(t) \cos(\theta(t)), r(t) \sin(\theta(t)), -f(t)) \\
z(t) &= (-r(t) \cos(\theta(t)), -r(t) \sin(\theta(t)), -f(t)),
\end{aligned}$$

is a solution of the 3-Body problem with the mass for the body moving according to $x(t)$ equal 200 and the masses of the other two bodies equal 100. We are assuming that the gravitational constant is 1.

Proof. Notice that $|x - y| = |x - z| = h$ and $|y - x| = 2r$, we can check that if f and r satisfy the ODE, then $\ddot{x}(t) = 100 \frac{y-x}{|y-x|^3} + 100 \frac{z-x}{|z-x|^3}$, $\ddot{y}(t) = 200 \frac{x-y}{|x-y|^3} + 100 \frac{z-y}{|z-y|^3}$ and $\ddot{z}(t) = 200 \frac{x-z}{|x-z|^3} + 100 \frac{y-z}{|y-z|^3}$. For more details see [6] or [3] for example. $\square$

Lemma 4.2. *Let us assume that f and g satisfies the ordinary differential equation*

$$\begin{aligned}
\ddot{f} &= \phi(f, g) \\
\ddot{g} &= \xi(f, g)
\end{aligned}$$

with $f(0) = 0$, $\dot{f}(0) = a$, $g(0) = r > 0$ and $\dot{g}(0) = 0$ and ϕ and ξ smooth functions. If $\phi(-f, g) = -\phi(f, g)$ and $\xi(-f, g) = \xi(f, g)$ then, $f(t)$ is odd and $g(t)$ is even.

Proof. Let us consider the functions $\tilde{f}(t) = -f(-t)$ and $\tilde{g}(t) = g(-t)$. A direct computation shows that $\tilde{f}(0) = 0$, $\dot{\tilde{f}}(0) = a$, $\tilde{g}(0) = r$ and $\dot{\tilde{g}}(0) = 0$. Moreover we have that

$$\ddot{\tilde{f}}(t) = -\ddot{f}(-t) = -\phi(f(-t), g(-t)) = -\phi(-\tilde{f}(t), \tilde{g}(t)) = \phi(\tilde{f}(t), \tilde{g}(t))$$

and

$$\ddot{\tilde{g}}(t) = \ddot{g}(-t) = \xi(f(-t), g(-t)) = \xi(-\tilde{f}(t), \tilde{g}(t)) = \xi(\tilde{f}(t), \tilde{g}(t))$$

By the uniqueness of the solutions of ordinary differential equations we get that $f(t) = \tilde{f}(t) = -f(-t)$ and $g(t) = \tilde{g}(t) = g(-t)$. This finishes the proof. $\square$

Lemma 4.3. *Let us assume that f and g satisfies the ordinary differential equation*

$$\begin{aligned}\ddot{f} &= \phi(f, g) \\ \ddot{g} &= \xi(f, g)\end{aligned}$$

If $\dot{f}(0) = 0$ and $\dot{g}(0) = 0$ and ϕ and ξ smooth functions, then, $f(t)$ and $g(t)$ are even.

Proof. Let us consider the functions $\tilde{f}(t) = f(-t)$ and $\tilde{g}(t) = g(-t)$. A direct computation shows that $\tilde{f}(0) = f(0)$, $\dot{\tilde{f}}(0) = 0$, $\tilde{g}(0) = g(0)$ and $\dot{\tilde{g}}(0) = 0$. Moreover we have that

$$\ddot{\tilde{f}}(t) = \ddot{f}(-t) = \phi(f(-t), g(-t)) = \phi(\tilde{f}(t), \tilde{g}(t))$$

and

$$\ddot{\tilde{g}}(t) = \ddot{g}(-t) = \xi(f(-t), g(-t)) = \xi(\tilde{f}(t), \tilde{g}(t))$$

By the uniqueness of the solutions of ordinary differential equations we get that $f(t) = \tilde{f}(t) = f(-t)$ and $g(t) = \tilde{g}(t) = g(-t)$. This finishes the proof. □

This is the Poincare-Miranda theorem for two variables

Theorem 4.4. *Let U be an open set in R^2 that contains the rectangle $[a_1, a_2] \times [t_1, t_2]$. Let us further assume that $F : U \times I \longrightarrow R$ and $G : U \times I \longrightarrow R$ are continuous functions. If*

(i) $F(a, t_1) > 0$ and $F(a, t_2) < 0$ for all $a \in [a_1, a_2]$,

(ii) $G(a_1, t) < 0$ and $G(a_2, t) > 0$ for all $t \in [t_1, t_2]$,

then, there exists a point $(a_0, t_0) \in [a_1, a_2] \times [t_1, t_2]$ such that $F(a_0, t_0) = 0 = G(a_0, t_0)$.

5 The solution of the ODE as a function of the time and the parameters a and b .

The main tool used to show the periodicity of the solution of the three body problem is to understand the functions in the solution of an ODE as functions of t , a and b .

We this in mind we define,

Definition 1. We will denote by $F(t, a, b) = f(t)$ and $R(t, a, b) = r(t)$ and $\Theta(t, a, b) = \theta(t)$ the solution of the system (4.1) with initial conditions

$$f(0) = 0, \quad \dot{f}(0) = b, \quad r(0) = 10, \quad \dot{r}(0) = 0 \quad \text{and} \quad \theta(0) = 0$$

We will denote by $\dot{R} = \frac{\partial R}{\partial t}$ and $\dot{F} = \frac{\partial F}{\partial t}$ and in general if S is a function of the variable (t, a, b) then $\dot{S} = \frac{\partial S}{\partial t}$

It is well know that the function F and R has continuous partial derivatives [2] and they obey a differential equation.

Theorem 5.1. If we denote by $F_a = \frac{\partial F}{\partial t}$, $F_b = \frac{\partial F}{\partial b}$, $R_a = \frac{\partial R}{\partial a}$ and $R_b = \frac{\partial R}{\partial b}$, then for any fixed values a and b ,

$$\begin{aligned} Y(t) = & (F(t, a, b), \dot{F}(t, a, b), R(t, a, b), \dot{R}(t, a, b), F_a(t, a, b), \dot{F}_a(t, a, b), \\ & R_a(t, a, b), \dot{R}_a(t, a, b), F_b(t, a, b), \dot{F}_b(t, a, b), R_b(t, a, b), \dot{R}_b(t, a, b)) \end{aligned}$$

satisfies the differential equation $\dot{Y} = P(Y)$ with $P(x_1, \dots, x_{12}) = P(x) = (P^1(x), \dots, P^{12}(x))$ where

$$\begin{aligned}
P^1(x) &= x_2 \\
P^2(x) &= -\frac{400x_1}{(4x_1^2 + x_3^2)^{3/2}} \\
P^3(x) &= x_4 \\
P^4(x) &= \frac{100a^2}{x_3^3} - \frac{25}{x_3^2} - \frac{200x_3}{(4x_1^2 + x_3^2)^{3/2}} \\
P^5(x) &= x_6 \\
P^6(x) &= \frac{3200x_5x_1^2}{(4x_1^2 + x_3^2)^{5/2}} + \frac{1200x_3x_7x_1}{(4x_1^2 + x_3^2)^{5/2}} - \frac{400x_3^2x_5}{(4x_1^2 + x_3^2)^{5/2}} \\
P^7(x) &= x_8 \\
P^8(x) &= -\frac{300a^2x_7}{x_3^4} + \frac{200a}{x_3^3} + \frac{2400x_1x_3x_5}{(4x_1^2 + x_3^2)^{5/2}} + \frac{50x_7}{x_3^3} + \frac{600x_3^2x_7}{(4x_1^2 + x_3^2)^{5/2}} - \frac{200x_7}{(4x_1^2 + x_3^2)^{3/2}} \\
P^9(x) &= x_{10} \\
P^{10}(x) &= \frac{3200x_9x_1^2}{(4x_1^2 + x_3^2)^{5/2}} + \frac{1200x_3x_{11}x_1}{(4x_1^2 + x_3^2)^{5/2}} - \frac{400x_3^2x_9}{(4x_1^2 + x_3^2)^{5/2}} \\
P^{11}(x) &= x_{12} \\
P^{12}(x) &= -\frac{300a^2x_{11}}{x_3^4} + \frac{2400x_1x_3x_9}{(4x_1^2 + x_3^2)^{5/2}} + \frac{50x_{11}}{x_3^3} + \frac{600x_3^2x_{11}}{(4x_1^2 + x_3^2)^{5/2}} - \frac{200x_{11}}{(4x_1^2 + x_3^2)^{3/2}}
\end{aligned}$$

with initial conditions

$$Y(0) = (0, b, 10, 0, 0, 0, 0, 0, 0, 1, 0, 0)$$

6 Main result

In this section we explain the arguments used in the proof of the periodicity of the solution and we will also show how to reduced this proof to the proof of 4 lemmas. With the intension of not cutting the flow of main ideas, we will prove these lemmas in a different section.

6.1 Special values used in the proof

before we continue explaining the main result we would like to defined some constants.

$$\begin{aligned} a_0 &= \frac{43170475352787}{10000000000000} & da &= \frac{17}{50000000} & sa &= \frac{1197}{100000000} \\ b_0 &= \frac{1490359743}{1000000000} & sb &= \frac{1}{50000} \\ t_0 &= \frac{13366894627923}{5000000000000} & dt &= \frac{1}{2500000} & st &= \frac{11}{2000000} \end{aligned}$$

6.2 Main Theorem: Periodic solution

Our main theorem shows that for some values of a , b and T of the form $a = 4.3170\dots$, $b = 1.4903\dots$ and $T = 96.24\dots$, the solution of the three body problem is periodic. For this periodic solution the orbit of the body with mass 200 goes up and down on the z -axis, and the other two bodies moves around the z -axis; the orbit of one of them is shown in Figure 0.1.

Theorem 6.1. *There exist a triple $(\bar{a}, \bar{b}, \bar{t})$ with $|\bar{a} - a_0| < sa + da$, $|\bar{b} - b_0| < sb$ and $|\bar{t} - t_0| < 6(ds + dt)$, such that the solution of the three body problem given by*

$$\begin{aligned} x(t) &= (0, 0, F(t, \bar{a}, \bar{b})) \\ y(t) &= (R(t, \bar{a}, \bar{b}) \cos(\Theta(t, \bar{a}, \bar{b})), R(t, \bar{a}, \bar{b}) \sin(\Theta(t, \bar{a}, \bar{b})), -F(t, \bar{a}, \bar{b})) \\ z(t) &= (-R(t, \bar{a}, \bar{b}) \cos(\Theta(t, \bar{a}, \bar{b})), -R(t, \bar{a}, \bar{b}) \sin(\Theta(t, \bar{a}, \bar{b})), -F(t, \bar{a}, \bar{b})), \end{aligned}$$

is period with period $36\bar{t}$. Moreover, we have that $\Theta(\bar{a}, \bar{b}, \bar{t}) = \frac{7\pi}{18}$.

6.3 Symmetry lemmas and reduced periodic solutions

A solution of the three body problem is called *reduced* periodic if the functions that provide the distances between the bodies are periodic with the same period T_0 . In the subfamily of solutions that we are describing in this paper, reduced periodic means that for some $T_0 > 0$, $R(t + kT_0, a, b) = R(t, a, b)$ and $f(t + kT_0, a, b) = f(t, a, b)$ for any integer k . The following theorem makes easier the task of finding reduced periodic solutions.

Lemma 6.2. *If for some $t_1 > 0$ we have that $\dot{R}(t_1, a, b) = 0 = \dot{F}(t_1, a, b)$, then the functions $f(t) = F(t, a, b)$ and $r(t) = R(t, a, b)$ are periodic with period $4t_1$.*

Proof. By Lemma (4.3) we have that $F(t_1 - t, a, b) = F(t_1 + t, a, b)$ and $R(t_1 - t, a, b) = R(t_1 + t, a, b)$. It follows that $-\dot{F}(t_1 - t, a, b) = \dot{F}(t_1 + t, a, b)$ and $-\dot{R}(t_1 - t, a, b) = \dot{R}(t_1 + t, a, b)$, therefore $\dot{R}(2t_1, a, b) = \dot{R}(0, a, b) = 0$ and $F(2t_1, a, b) = F(0, a, b) = 0$. By Lemma 4.2 we have that $f(-t) = -f(t)$ and $r(-t) = r(t)$. Using Lemma 4.2 one more time we obtain that $f(t + 2t_1) = -f(2t_1 - t)$ and $r(2t_1 + t) = r(2t_1 - t)$. It follows that $f(4t_1) = f(0) = 0$, $\dot{f}(4t_1) = \dot{f}(0) = b$, $r(4t_1) = r(0) = 10$ and $\dot{r}(4t_1) = -\dot{r}(0) = 0$. Since the values of the solutions f and r at $t = 4t_1$ agrees with those at $t = 0$, then the lemma follows. $\square$

By the previous theorem, a point (t_1, a_1, b_1) satisfying $\dot{R}(t_1, a_1, b_1) = 0 = \dot{F}(t_1, a_1, b_1)$ defines a reduced periodic solution. If additionally $\Theta(t_1, a_1, b_1) = \frac{2\pi p}{q}$ with p and $q > 0$ integers, then the solution is not only reduced periodic but periodic with period $4qt_1$. Moreover, by the implicit function theorem if the cross product between the gradients of the functions $\dot{R}$ and $\dot{F}$ at (t_1, a_1, b_1) does not vanish, then there is a curve of points that solve the equation $\dot{R}(t, a, b) = 0 = \dot{F}(t, a, b)$ and then, we obtain a family of reduced periodic solutions. In the case that the function Θ is not constant along this curve of points in the space that represent reduced periodic solutions, then we obtain infinitely many periodic solution due to the fact that on any open interval there are infinitely many numbers of the form $\frac{2\pi p}{q}$ with p and q integers. The Implicit function theorem also tell us that there exists an small open set around (t_1, a_1, b_1) such that all the solutions of the equations $\dot{R}(t, a, b) = 0 = \dot{F}(t, a, b)$ in this small open set must be part of this curve... but how small is small? Notice that the following two difficulties need to be taken care of: (i) The fact that $\Theta(t_1, a_1, b_1)$ is near $\frac{7\pi}{18}$ and Θ is not constant along the curve, does not imply that Θ eventually reaches the value $\frac{7\pi}{18}$ on this curve. (ii) The fact that $\Theta(t_1, a_1, b_1) < \frac{7\pi}{18}$ and $\Theta(t_2, a_2, b_2) > \frac{7\pi}{18}$ for two nearby points (t_1, a_1, b_1) and (t_2, a_2, b_2) that satisfy the equations $\dot{R}(t, a, b) = 0 = \dot{F}(t, a, b)$, does not guarantee that Θ eventually reaches the value $\frac{7\pi}{18}$ due to the fact the point (t_1, a_1, b_1) may no be in the same connected component of the curve of solutions that contains (t_2, a_2, b_2) . Theorem 3.1 helps to solve these two difficulties.

The idea of the proof of Theorem 6.1 is the following. We consider a box in the space, see Figure 6.1. We use Theorem 3.1 to show that every plane $b = c$ in this box contains only one solution of the equation $\dot{R}(t, a, b) = 0 = \dot{F}(t, a, b)$. We use the Poincare Miranda Theorem to show that there are reduced periodic solutions on three rectangles contained in planes of the form $b = b_0$, $b = b_0 - sb$ and $b = b_0 + sb$. We finish the prove by showing that Θ evaluated on the solution of the equations




Figure 6.1: Points (t, a, b) in the curve represent reduced periodic solutions of the three body problem. We proved the existence of a point in this curve in each one of the three small rectangles shown in the picture and we prove that at the point in the curve contained in the first rectangle the value of the function Θ is smaller than $\frac{7\pi}{18}$ and at the point in the curve contained in the last rectangle the value of the function Θ is bigger than $\frac{7\pi}{18}$. The proof of the existence of a point in the curve on the rectangle in the center was needed to implement our theorem regarding the Implicit Function Theorem, which, in this case, it guarantees the existence of a unique point on the curve in every plane $b = c$.

$\dot{R}(t, a, b) = 0 = \dot{F}(t, a, b)$ contained on the rectangle with $b = b_0 - sb$ is less than $\frac{7\pi}{18}$ and Θ evaluated on the solution of the equations $\dot{R}(t, a, b) = 0 = \dot{F}(t, a, b)$ contained on the rectangle with $b = b_0 + sb$ is greater than $\frac{7\pi}{18}$.

Now we are ready to state the four lemmas mentioned in the beginning of this section that lead to the proof of the main Theorem 6.1. The first lemma shows that there is a point in the curve on the box in Figure 6.1 in the rectangle in the center.

Lemma 6.3. *There is a solution of the equations $\dot{R}(t, a, b) = 0 = \dot{F}(t, a, b)$ with $b = b_0$, $|a - a_0| < da$ and $|t - t_0| < dt$*

The second and third lemmas show that there is a point in the curve on the initial and final rectangles 6.1.

Lemma 6.4. *For some (t, a, b) with $b = b_0 - sb$, $|a - (a_0 + sa)| < da$ and $|t - (t_0 - st)| < dt$, there is a solution of the equations $\dot{R}(t, a, b) = 0 = \dot{F}(t, a, b)$ such that $\Theta(t, a, b) < \frac{7\pi}{18}$.*

Lemma 6.5. *For some (t, a, b) with $b = b_0 + sb$, $|a - (a_0 - sa)| < da$ and $|t - (t_0 + st)| < dt$, there is a solution of the equations $\dot{R}(t, a, b) = 0 = \dot{F}(t, a, b)$ such that $\Theta(t, a, b) > \frac{7\pi}{18}$.*

The fourth lemma guaranties the set for values of (t, a, b) given by

$$[t_0 - 6(st + dt), t_0 + 6(st + dt)] \times [a_0 - 3(sa + da), a_0 + 3(sa + da)] \times [b_0 - sb, b_0 + sb]$$

is small enough to only allow one connected curve on it as the solution for the equation $\dot{R}(t, a, b) = 0 = \dot{F}(t, a, b)$.

Lemma 6.6. *For every $b \in [b_0 - sb, b_0 + sb]$ there exists a unique solution (t, a, b) of the equations $\dot{R}(t, a, b) = 0 = \dot{F}(t, a, b)$ with $|a - a_0| < 3(sa + da)$ and $|t - t_0| < 6(st + dt)$.*

7 Bounds

In this section we will define 3 differential equations that will help us find bounds for the functions $F(t, a, b)$, $R(t, a, b)$, $\Theta(t, a, b)$ and their partial derivatives. We will be using the functions ϕ_i defined in section 8.

7.1 The vector field W

Let us consider the vector field $W = (x_2, \phi_1, x_4, \phi_2, \phi_3)^T$, where the functions ϕ_i 's are those defined on section 8. For any fixed a and b , the function

$$t \rightarrow \left(F(t, a, b), \dot{F}(t, a, b), R(t, a, b), \dot{R}(t, a, b), \Theta(t, a, b) \right)^T$$

satisfies the differential equation $Y(t) = W(Y(t))$ with $Y(0) = (0, b, 10, 0, 0)^T$

Proposition 7.1. *Using the functions ϕ_i defined in section 8, we have that the derivative matrix DW of the vector field W , and the derivative matrix DW_1 of the vector field $W_1 = DW W$ are given by,*

$$DW = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ \phi_4 & 0 & \phi_5 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 2\phi_5 & 0 & \phi_6 & 0 & 0 \\ 0 & 0 & \phi_7 & 0 & 0 \end{pmatrix} \quad \text{and} \quad DW_1 = \begin{pmatrix} \phi_4 & 0 & \phi_5 & 0 & 0 \\ \phi_{10} & \phi_4 & \phi_{12} & \phi_5 & 0 \\ 2\phi_5 & 0 & \phi_6 & 0 & 0 \\ 2\phi_{12} & 2\phi_5 & \phi_{14} & \phi_6 & 0 \\ 0 & 0 & \phi_{15} & \phi_7 & 0 \end{pmatrix}$$

7.2 The vector field G

Let us consider the vector field $G = (x_2, \phi_1, x_4, \phi_2, x_6, \phi_{16}, x_8, \phi_{17}, \phi_{18})^T$, where the function ϕ_i are those defined on section 8. For any fixed a and b , the function that sends t to

$$\left(F(t, a, b), \dot{F}(t, a, b), R(t, a, b), \dot{R}(t, a, b), F_a(t, a, b), \dot{F}_a(t, a, b), R_a(t, a, b), \dot{R}_a(t, a, b), \Theta_a(t, a, b) \right)^T$$

satisfies the differential equation $Y(t) = G(Y(t))$ with $Y(0) = (0, b, 10, 0, 0, 0, 0, 0, 0)^T$

Proposition 7.2. *Using the functions ϕ_i defined on section 8, we have that the derivative matrix DG of the vector field G , and the derivative matrix DG_1 of the vector field $G_1 = DG G$ are given by,*

$$DG = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \phi_4 & 0 & \phi_5 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 2\phi_5 & 0 & \phi_6 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ \phi_{19} & 0 & \phi_{20} & 0 & \phi_4 & 0 & \phi_5 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 2\phi_{20} & 0 & \phi_{22} & 0 & 2\phi_5 & 0 & \phi_6 & 0 & 0 \\ 0 & 0 & \phi_{23} & 0 & 0 & 0 & \phi_7 & 0 & 0 \end{pmatrix}$$

and

$$DG_1 = \begin{pmatrix} \phi_4 & 0 & \phi_5 & 0 & 0 & 0 & 0 & 0 & 0 \\ \phi_{10} & \phi_4 & \phi_{12} & \phi_5 & 0 & 0 & 0 & 0 & 0 \\ 2\phi_5 & 0 & \phi_6 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2\phi_{12} & 2\phi_5 & \phi_{14} & \phi_6 & 0 & 0 & 0 & 0 & 0 \\ \phi_{19} & 0 & \phi_{20} & 0 & \phi_4 & 0 & \phi_5 & 0 & 0 \\ \phi_{30} & \phi_{19} & \phi_{34} & \phi_{20} & \phi_{10} & \phi_4 & \phi_{12} & \phi_5 & 0 \\ 2\phi_{20} & 0 & \phi_{22} & 0 & 2\phi_5 & 0 & \phi_6 & 0 & 0 \\ 2\phi_{34} & 2\phi_{20} & \phi_{39} & \phi_{22} & 2\phi_{12} & 2\phi_5 & \phi_{14} & \phi_6 & 0 \\ 0 & 0 & \phi_{40} & \phi_{23} & 0 & 0 & \phi_{15} & \phi_7 & 0 \end{pmatrix}$$

7.3 The vector field U

Let us consider the vector field

$$U = (x_2, \phi_1, x_4, \phi_2, x_{10}, \phi_{41}, x_{12}, \phi_{42})^T,$$

where the function ϕ_i are those defined on section 8. For any fixed a and b the function

$$t \rightarrow \begin{pmatrix} F(t, a, b), \dot{F}(t, a, b), R(t, a, b), \dot{R}(t, a, b), F_b(t, a, b), \dot{F}_b(t, a, b), \\ R_b(t, a, b), \dot{R}_b(t, a, b) \end{pmatrix}^T$$

satisfies the differential equation $\dot{Y}(t) = U(Y(t))$ with $Y(0) = (0, b, 10, 0, 0, 1, 0, 0)^T$

Proposition 7.3. *Using the functions ϕ_i defined in section 8, we have that the derivative matrix DU of the vector field U , and the derivative matrix DU_1 of the vector field $U_1 = DUU$ are given by,*

$$DU = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ \phi_4 & 0 & \phi_5 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 2\phi_5 & 0 & \phi_6 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ \phi_{43} & 0 & \phi_{44} & 0 & \phi_4 & 0 & \phi_5 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 2\phi_{44} & 0 & \phi_{45} & 0 & 2\phi_5 & 0 & \phi_6 & 0 \end{pmatrix}$$

and

$$DU_1 = \begin{pmatrix} \phi_4 & 0 & \phi_5 & 0 & 0 & 0 & 0 & 0 \\ \phi_{10} & \phi_4 & \phi_{12} & \phi_5 & 0 & 0 & 0 & 0 \\ 2\phi_5 & 0 & \phi_6 & 0 & 0 & 0 & 0 & 0 \\ 2\phi_{12} & 2\phi_5 & \phi_{14} & \phi_6 & 0 & 0 & 0 & 0 \\ \phi_{43} & 0 & \phi_{44} & 0 & \phi_4 & 0 & \phi_5 & 0 \\ \phi_{48} & \phi_{43} & \phi_{50} & \phi_{44} & \phi_{10} & \phi_4 & \phi_{12} & \phi_5 \\ 2\phi_{44} & 0 & \phi_{45} & 0 & 2\phi_5 & 0 & \phi_6 & 0 \\ 2\phi_{50} & 2\phi_{44} & \phi_{53} & \phi_{45} & 2\phi_{12} & 2\phi_5 & \phi_{14} & \phi_6 \end{pmatrix}$$

7.4 Reduced Periodic Solutions

In this section we use theorem 2.3 and the Poincare -Miranda Theorem to prove the existence of three reduced periodic solutions, one on each rectangle in figure 6.1.

Remark 7.4. *Several lemmas in this section will be using the Round Taylor method and therefore it will be using Theorem 2.3 to estimate the values of the solution of the ODE's. If we take a look at the hypothesis of this lemma we notice that there is a number ϵ that has to be greater than $M_0h + \tilde{H}$. This ϵ will be $\frac{1}{1000}$ in all proofs that use the Taylor method in this paper. Some of these lemmas contains the variable ϵ , in each case, it just refers to a small number giving an estimate of the error.*

Lemma 7.5. *Let $\epsilon = \frac{2677451}{100000000}$. For any $a \in [a_0 - 3(sa + da), a_0 + 3(sa + da)]$, $b \in [b_0 - sb, b_0 + sb]$ and $t \in [t_0 - 6(st + dt), t_0 + 6(st + dt)]$ we have that*

$$|\Theta_a(t, a, b) - \frac{536760312951}{20000000000000}| < \epsilon$$

and,

$$|F_a(t, a, b) - \frac{3032500537707}{10000000000000}| < \epsilon, \quad |\dot{F}_a(t, a, b) - \frac{11824770099363}{25000000000000}| < \epsilon$$

$$|R_a(t, a, b) - \frac{68073031375453}{25000000000000}| < \epsilon, \quad |\dot{R}_a(t, a, b) - \frac{164497338366219}{100000000000000}| < \epsilon$$

Proof. Let us consider the following intervals,

$$I_1 = \left[-\frac{1}{100}, \frac{62}{25}\right] \quad I_2 = \left[-\frac{1}{100}, \frac{3}{2}\right] \quad I_3 = \left[\frac{189}{20}, \frac{1001}{100}\right] \quad I_4 = \left[-\frac{33}{100}, \frac{1}{100}\right] \quad I_5 = \left[-\frac{1}{100}, \frac{31}{100}\right]$$

$$I_6 = \left[-\frac{1}{100}, \frac{12}{25}\right] \quad I_7 = \left[-\frac{1}{100}, \frac{69}{25}\right] \quad I_8 = \left[-\frac{1}{100}, \frac{42}{25}\right] \quad J_9 = \left[-\frac{1}{100}, \frac{3}{25}\right]$$

A direct computation using the bounds in section 8 shows that if

$$M_{0g} = \frac{42}{25}, \quad K_{0g} = \frac{305541}{125000}, \quad K_{1g} = \frac{249309}{100000}, \quad M_{1g} = \frac{778131}{1000000} \quad M_{2g} = \frac{246743}{250000}$$

$$M_{3g} = \frac{374443}{500000}, \quad M_{4g} = \frac{133409}{100000}, \quad M_{5g} = \frac{1345793}{1000000}, \quad M_{6g} = \frac{2429239}{1000000}$$

$$M_{7g} = \frac{833241}{500000}, \quad M_{8g} = \frac{3298559}{1000000}, \quad M_9 = \frac{182893}{1000000}, \quad H = 10^{-14}, \quad h = \frac{t_0}{30000},$$

then, for values of $(x_1, \dots, x_8)$ with $x_i \in I_i$ we have that $|G^j| < M_{0g}$ where $G^1, \dots, G^9$ are the entries of the vector field G ; $|G_2^j| < M_{jg}$ where $G_2^1, \dots, G_2^9$ are the entries of the vector field $G_2 = DG_1 G$. Recall that $G_1 = DG G$. Moreover we have that

$$|DG| < K_{0g}, \quad |DG_1| < K_{1g}$$

The Round Taylor method of order 2 using the vector field G with $a = a_0$, $k = 30000$ and initial conditions $Y(0) = (0, b_0, 10, 0, 0, 0, 0, 0)^T$ produces a sequence $\{z_i\}_{i=1}^k$ with z_{30000} equal to

$$\left(\frac{247458249564811}{100000000000000}, \frac{13245901}{100000000000000}, \frac{189061430242601}{200000000000000}, \frac{1795639}{125000000000000}, \right. \\ \left. \frac{3032500537707}{100000000000000}, \frac{11824770099363}{250000000000000}, \frac{68073031375453}{250000000000000}, \frac{164497338366219}{100000000000000}, \frac{536760312951}{200000000000000} \right)$$

In this case $\tilde{H} = \frac{M_g \frac{h^2}{6} + \frac{H}{h}}{L} (e^{Lkh} - 1) \approx 0.0000019$ where $M_g = \sqrt{M_{1g}^2 + \dots + M_{9g}^2}$ and $L = K_{0g} + K_{1g} \frac{h}{2}$. A direct verification shows that, for every $j = 1, \dots, 8$ and $i = 1, \dots, 30000$, the j^{th} entry of z_i is within a distance ϵ of the boundary of the interval I_j . Also we have that the last entry of z_i is within a distance $\frac{1}{1000}$ of the boundary of the interval J_9 . By Theorem 2.3 we conclude that for all $i = 1, \dots, k$,

$$\left(F\left(\frac{ih}{k}, a_0, b_0\right), \dot{F}\left(\frac{ih}{k}, a_0, b_0\right), R\left(\frac{ih}{k}, a_0, b_0\right), \dot{R}\left(\frac{ih}{k}, a_0, b_0\right), F_a\left(\frac{ih}{k}, a_0, b_0\right), \dot{F}_a\left(\frac{ih}{k}, a_0, b_0\right), \right. \\ \left. R_a\left(\frac{ih}{k}, a_0, b_0\right), \dot{R}_a\left(\frac{ih}{k}, a_0, b_0\right), \Theta_a\left(\frac{ih}{k}, a_0, b_0\right) \right)^T$$

is within a distance $\tilde{H}$ of z_i . Notice that for any function ρ , under the assumption that $|\dot{\rho}(\tau, a_0, b_0)| < M_\rho$ for all τ between t and t_0 , we have that

$$\begin{aligned} |\rho(t, a, b) - \rho(t_0, a_0, b_0)| &\leq |\rho(t, a, b) - \rho(t, a_0, b_0)| + |\rho(t, a_0, b_0) - \rho(t_0, a_0, b_0)| \\ &\leq |\rho(t, a, b) - \rho(t, a_0, b_0)| + M_\rho (t - t_0), \end{aligned}$$

We will use the observation above to finish the proof of the lemma. We will bound $|\rho(t, a, b) - \rho(t, a_0, b_0)|$ using Theorem 2.4 with the vector fields G and G_0 , where G_0 is the vector field G with a replaced by a_0 . A direct computation shows that

$$\delta G = G - G_0 = (0, 0, 0, \phi_{54}, 0, 0, 0, \phi_{56}, \phi_{57})$$

Using the information on section 8 we obtain that $|\delta G| < \frac{23}{500000}$. Therefore, using Theorem 2.4 we conclude that the values of the solution of the differential equation using G (with a general a and b) compare with those of the solution of the differential equation using G_0 differ by less than

$$|b - b_0| e^{K_{0g}(t_0 + 6(st + dt))} + \frac{23}{500000 K_{0g}} \left(e^{K_{0g}(t_0 + 6(st + dt))} - 1 \right) < \frac{267131}{10000000}$$

Therefore we have that for any $a \in [a_0 - 3(sa + da), a_0 + 3(sa + da)]$, $b \in [b_0 - sb, b_0 + sb]$ and $t \in [t_0 - 6(st + dt), t_0 + 6(st + dt)]$

$$|F_a(t, a, b) - \frac{3032500537707}{10000000000000}| < \frac{267131}{10000000} + 6(st + dt)M_{0g} + \tilde{H} \leq \frac{2677451}{100000000}$$

We have similar computations for the functions Θ_a , $\dot{F}_a$, R_a and $\dot{R}_a$. This finishes the proof. $\square$

Corollary 7.6. *For any $a \in [a_0 - 3(sa + da), a_0 + 3(sa + da)]$, $b \in [b_0 - sb, b_0 + sb]$ and $t \in [t_0 - 6(st + dt), t_0 + 6(st + dt)]$, $\dot{F}_a(t, a, b) > 0$, $\dot{R}_a(t, a, b) > 0$, and $|\Theta_a(t, a, b)| < \frac{27}{1000}$*

Lemma 7.7. *Let $\epsilon = \frac{2568201}{1000000000}$. For any $a \in [a_0 - 3(sa + da), a_0 + 3(sa + da)]$, $b \in [b_0 - sb, b_0 + sb]$ and $t \in [t_0 - 6(st + dt), t_0 + 6(st + dt)]$ we have that*

$$|\dot{R}_b(t, a, b) - \frac{88229751956717}{100000000000000}| < \epsilon, \quad |\dot{F}_b(t, a, b) - \frac{50798112898451}{100000000000000}| < \epsilon$$

Proof. Let us consider the following intervals,

$$I_1 = \left[-\frac{1}{100}, \frac{62}{25}\right] \quad I_2 = \left[-\frac{1}{100}, \frac{3}{2}\right] \quad I_3 = \left[\frac{189}{20}, \frac{1001}{100}\right] \quad I_4 = \left[-\frac{33}{100}, \frac{1}{100}\right]$$

$$I_9 = \left[-\frac{1}{100}, \frac{101}{50}\right] \quad I_{10} = \left[\frac{1}{2}, \frac{101}{100}\right] \quad I_{11} = \left[-\frac{1}{100}, \frac{81}{100}\right] \quad I_{12} = \left[-\frac{1}{100}, \frac{89}{100}\right]$$

A direct computation using section 8 shows that if

$$M_{0u} = \frac{3}{2}, \quad K_{0u} = \frac{1226931}{500000}, \quad K_{1u} = \frac{2557349}{1000000}, \quad M_{1u} = \frac{778131}{1000000}, \quad M_{2u} = \frac{246743}{250000}$$

$$M_{3u} = \frac{374443}{500000}, \quad M_{4u} = \frac{133409}{100000}, \quad M_{5u} = \frac{765259}{500000}, \quad M_{6u} = \frac{533571}{200000}$$

$$M_{7u} = \frac{1790753}{1000000}, \quad M_{8u} = \frac{711267}{200000}, \quad H = 10^{-14}, \quad h = \frac{t_0}{30000},$$

then, for values of $(x_1, x_2, x_3, x_4, x_9, x_{10}, x_{11}, x_{12})$ with $x_i \in I_i$ we have that $|U^j| < M_{0g}$ where $U^1, \dots, U^8$ are the entries of the vector field U ; $|U_2^j| < M_{ju}$ where

$U_2^1, \dots, U_2^8$ are the entries of the vector field $U_2 = DU_1 U$. Recall that $U_1 = DU U$. Moreover we have that

$$|DU| < K_{0u}, \quad |DU_1| < K_{1u}$$

The Round Taylor method of order 2 using the vector field U with $a = a_0$, $k = 30000$ and initial conditions $Y(0) = (0, b_0, 10, 0, 0, 1, 0, 0)^T$ produces a sequence $\{z_i\}_{i=1}^k$ with z_{30000} equal to

$$\left(\frac{247458249564811}{10000000000000}, \frac{13245901}{10000000000000}, \frac{189061430242601}{20000000000000}, \frac{1795639}{12500000000000}, \right. \\ \left. \frac{25138479462137}{12500000000000}, \frac{50798112898451}{10000000000000}, \frac{20014508374143}{25000000000000}, \frac{88229751956717}{10000000000000} \right)$$

In this case $\tilde{H} = \frac{M_u \frac{h^2}{6} + \frac{H}{h}}{L} (e^{Lkh} - 1) < \frac{209}{1000000000}$ where $M_u = \sqrt{M_{1u}^2 + \dots + M_{8u}^2}$ and $L = K_{0u} + K_{1u} \frac{h}{2}$. A direct verification shows that, for every $j = 1, \dots, 8$ and $i = 1, \dots, 30000$, the j^{th} entry of z_i is within a distance $\frac{1}{1000}$ of the boundary of the interval I_j when $j = 1, 2, 3, 4$ and the j^{th} entry of z_i is within a distance $\frac{1}{1000}$ of the boundary of the interval I_{j+4} when $j = 5, 6, 7, 8$. By Theorem 2.3 we conclude that for all $i = 1, \dots, k$,

$$\left(F\left(\frac{ih}{k}, a_0, b_0\right), \dot{F}\left(\frac{ih}{k}, a_0, b_0\right), R\left(\frac{ih}{k}, a_0, b_0\right), \dot{R}\left(\frac{ih}{k}, a_0, b_0\right), F_b\left(\frac{ih}{k}, a_0, b_0\right), \dot{F}_b\left(\frac{ih}{k}, a_0, b_0\right), \right. \\ \left. R_b\left(\frac{ih}{k}, a_0, b_0\right), \dot{R}_b\left(\frac{ih}{k}, a_0, b_0\right) \right)^T$$

is within a distance $\tilde{H}$ of z_i . Notice that for any function ρ , under the assumption that $|\dot{\rho}(\tau, a_0, b_0)| < M_\rho$ for all τ between t and t_0 , we have that

$$\begin{aligned} |\rho(t, a, b) - \rho(t_0, a_0, b_0)| &\leq |\rho(t, a, b) - \rho(t, a_0, b_0)| + |\rho(t, a_0, b_0) - \rho(t_0, a_0, b_0)| \\ &\leq |\rho(t, a, b) - \rho(t, a_0, b_0)| + M_\rho (t - t_0), \end{aligned}$$

We will use the observation above to finish the proof of the lemma. We will bound $|\rho(t, a, b) - \rho(t, a_0, b_0)|$ using Theorem 2.4 with the vector fields U and U_0 , where U_0 is the vector field U with a replaced by a_0 . A direct computation shows that

$$\delta U = U - U_0 = (0, 0, 0, \phi_{54}, 0, 0, 0, \phi_{58})$$

Using the information on section 8 we obtain that $|\delta_U| < \frac{1}{25000}$. Therefore, using Theorem 2.4 we conclude that the values of the the solution of the differential equation using U (with a general a and b) compare with those of the solution of the differential equation using U_0 differ by less than

$$|b - b_0|e^{K_{0u}(t_0+6(st+dt))} + \frac{1}{25000K_{0u}} \left(e^{K_{0u}(t_0+6(st+dt))} - 1 \right) < \frac{1281341}{50000000}$$

Therefore we have that for any $a \in [a_0 - 3(sa+da), a_0 + 3(sa+da)]$, $b \in [b_0 - sb, b_0 + sb]$ and $t \in [t_0 - 6(st+dt), t_0 + 6(st+dt)]$

$$|\dot{F}_b(t, a, b) - \frac{50798112898451}{100000000000000}| < \frac{1281341}{50000000} + 6(st+dt)M_{0u} + \tilde{H} \leq \frac{2568201}{100000000}$$

We have a similar computation for the function $\dot{R}_b$. This finishes the proof. $\square$

Lemma 7.8. *Let $\epsilon = \frac{134567}{40000000}$. For any $a \in [a_0 - 3(sa+da), a_0 + 3(sa+da)]$, $b \in [b_0 - sb, b_0 + sb]$ and $t \in [t_0 - 6(st+dt), t_0 + 6(st+dt)]$ we have that*

$$|F(t, a, b) - \frac{247458249564811}{100000000000000}| < \epsilon, \quad |R(t, a, b) - \frac{189061430242601}{20000000000000}| < \epsilon$$

Proof. Let us consider the following intervals $J_5 = [-\frac{1}{100}, \frac{7}{5}]$ and I_1, I_2, I_3, I_4 defined on the proof of Lemma 7.5.

A direct computation using the bounds on section 8 shows that if

$$M_{0w} = \frac{3}{2}, \quad K_{0w} = \frac{25282}{15625}, \quad K_{1w} = \frac{260901}{200000}, \quad M_{1w} = \frac{19453263}{25000000}, \quad M_{3w} = \frac{14977713}{20000000}$$

$$M_{2w} = \frac{493485626283}{500000000000}, \quad M_{4w} = \frac{1334089805457}{1000000000000}, \quad M_{5w} = \frac{5396988231}{125000000000},$$

then, for values of $(x_1, \dots, x_4)$ with $x_i \in I_i$ we have that $|W^j| < M_{0w}$ where $W^1, \dots, W^5$ are the entries of the vector field W ; $|W_2^j| < M_{jw}$ where $W_2^1, \dots, W_2^5$ are the entries of the vector field $W_2 = DW_1 W$. Recall that $W_1 = DW W$. Moreover we have that

$$|DW| < K_{0w}, \quad |DW_1| < K_{1w}$$

The Round Taylor method of order 2 using the vector field W with $H = 10^{-14}$, $h = \frac{t_0}{30000}$, $a = a_0$, $k = 30000$ and initial conditions $Y(0) = (0, b_0, 10, 0, 0)^T$ produces a sequence $\{z_i\}_{i=1}^k$ with z_{30000} equal to

$$\left(\frac{247458249564811}{100000000000000}, \frac{13245901}{100000000000000}, \frac{189061430242601}{200000000000000}, \frac{1795639}{125000000000000}, \frac{12217304404331}{100000000000000} \right)$$

In this case $\tilde{H} = \frac{M_w \frac{h^2}{6} + \frac{H}{h}}{L} (e^{Lkh} - 1) < \frac{127}{1000000000}$ where $M_w = \sqrt{M_{1w}^2 + \dots + M_{5w}^2}$ and $L = K_{0w} + K_{1w} \frac{h}{2}$. A direct verification shows that, for every $j = 1, \dots, 4$ and $i = 1, \dots, 30000$, the j^{th} entry of z_i is within a distance $\frac{1}{1000}$ of the boundary of the interval I_j . Also we have that the last entry of z_i is within a distance $\frac{1}{1000}$ of the boundary of the interval J_5 . By Theorem 2.3 we conclude that for all $i = 1, \dots, k$,

$$\left(F\left(\frac{ih}{k}, a_0, b_0\right), \dot{F}\left(\frac{ih}{k}, a_0, b_0\right), R\left(\frac{ih}{k}, a_0, b_0\right), \dot{R}\left(\frac{ih}{k}, a_0, b_0\right), \Theta_a\left(\frac{ih}{k}, a_0, b_0\right) \right)^T$$

is within a distance $\tilde{H}$ of z_i . Notice that for any function ρ , under the assumption that $|\dot{\rho}(\tau, a_0, b_0)| < M_\rho$ for all τ between t and t_0 , we have that

$$\begin{aligned} |\rho(t, a, b) - \rho(t_0, a_0, b_0)| &\leq |\rho(t, a, b) - \rho(t, a_0, b_0)| + |\rho(t, a_0, b_0) - \rho(t_0, a_0, b_0)| \\ &\leq |\rho(t, a, b) - \rho(t, a_0, b_0)| + M_\rho(t - t_0), \end{aligned}$$

We will use the observation above to finish the proof of the lemma. We will bound $|\rho(t, a, b) - \rho(t, a_0, b_0)|$ using Theorem 2.4 with the vector fields W and W_0 , where W_0 is the vector field W with a replaced by a_0 . A direct computation shows that

$$\delta W = W - W_0 = (0, 0, 0, \phi_{54}, \phi_{55})$$

Using the information on section 8 we obtain that $|\delta W| < \frac{39}{1000000}$. Therefore, using Theorem 2.4 we conclude that the values of the the solution of the differential equation using W (with a general a and b) compare with those of the solution of the differential equation using W_0 differ by less than

$$|b - b_0| e^{K_{0w}(t_0 + 6(st + dt))} + \frac{39}{1000000 K_{0w}} \left(e^{K_{0w}(t_0 + 6(st + dt))} - 1 \right) < \frac{827737}{250000000}$$

Therefore we have that for any $a \in [a_0 - 3(sa + da), a_0 + 3(sa + da)]$, $b \in [b_0 - sb, b_0 + sb]$ and $t \in [t_0 - 6(st + dt), t_0 + 6(st + dt)]$

$$|F(t, a, b) - \frac{247458249564811}{100000000000000}| < \frac{827737}{250000000} + 6(st + dt)M_{0w} + \tilde{H} \leq \frac{134567}{40000000}$$

We have a similar computation for the function R . This finishes the proof. $\square$

Definition 2. We will denote by $Z_W(t, b, a, k) \in R^5$ the last vector in the sequence $\{z_i\}_{i=1}^k$ produced using the Round Taylor Method of order 2 using the vector field W with $H = 10^{-14}$, $h = \frac{t}{k}$ and initial conditions $(0, b, 10, 0, 0)^T$. We denote by $H_W(t, k) = \frac{M_w \frac{h^2}{6} + \frac{H}{h}}{L} (e^{Lkh} - 1)$ where $M_w = \sqrt{M_{1w}^2 + \dots + M_{5w}^2}$ and $L = K_{0w} + K_{1w} \frac{h}{2}$. If we need to use $H = 10^{-q}$ instead of $H = 10^{-14}$, then we will use the notation $Z_W(t, b, a, k, q) \in R^5$ and $H_W(t, k, q)$.

Remark 7.9. For all the following Lemmas that use the Round Taylor Method, it can be directly verified that all the conditions of Theorem 2.3 are satisfied; therefore $Z_W(t, b, a, k)$ is within a distance $H_W(t, k)$ of

$$(F(t, a, b), \dot{F}(t, a, b), R(t, a, b), \dot{R}(t, a, b), \Theta(t, a, b))^T.$$

For the sake of keeping the flow of the proof, we will left out all the details in the proofs of these lemmas.

Corollary 7.10. For any $a \in [a_0 - 3(sa + da), a_0 + 3(sa + da)]$, $b \in [b_0 - sb, b_0 + sb]$ and $t \in [t_0 - 6(st + dt), t_0 + 6(st + dt)]$ we have that $\frac{120689}{250000} < \dot{\Theta}(t, a, b) < \frac{483453}{1000000}$,

$$-\frac{163169}{200000} < \ddot{F}(t, a, b) < -\frac{813693}{1000000} \quad \text{and} \quad \frac{4593}{12500} < \ddot{R}(t, a, b) < \frac{18653}{50000}$$

Proof. As in Lemma 7.8, let $\epsilon = \frac{134567}{40000000}$. Notice that the values of $\ddot{F}(t, a, b)$ are bounded by the minimum and maximum of the function $-\frac{400x_1}{(4x_1^2 + x_3^2)^{3/2}}$ when $|x_1 - \frac{247458249564811}{100000000000000}| < \epsilon$ and $|x_3 - \frac{189061430242601}{200000000000000}| < \epsilon$. The bound for $\ddot{F}$ follows as a direct application of the Lagrange Multiplier method. Since the values of $\ddot{R}(t, a, b)$ are bounded by the minimum and maximum of the function $\frac{100a^2}{x_3^3} - \frac{25}{x_3^2} - \frac{200x_3}{(4x_1^2 + x_3^2)^{3/2}}$, we can use the same argument for $\ddot{R}(t, a, b)$. Finally since the values of $\dot{\Theta}(t, a, b)$ are bounded by the minimum and maximum of the function $\frac{10a}{x_3^2}$, we can also use the same argument for $\dot{\Theta}(t, a, b)$. $\square$

Lemma 7.11. $\dot{F}(t_0 - dt, a, b_0) > 0$ for all $a \in [a_0 - da, a_0 + da]$, $\dot{F}(t_0 + dt, a, b_0) < 0$ for all $a \in [a_0 - da, a_0 + da]$, $\dot{R}(t, a_0 - da, b_0) < 0$ for all $t \in [t_0 - dt, t_0 + dt]$ and $\dot{R}(t, a_0 + da, b_0) > 0$ for all $t \in [t_0 - dt, t_0 + dt]$.

Proof. By Corollary 7.6 we have that $\dot{F}_a(t_0 - dt, a, b_0) > 0$ for all $a \in [a_0 - da, a_0 + da]$. Therefore, in order to show that $\dot{F}(t_0 - dt, a, b_0) > 0$ for all $a \in [a_0 - da, a_0 + da]$, it is enough to show that $\dot{F}(t_0 - dt, a_0 - da, b_0) > 0$. A direct computation shows that $Z_W(t_0 - dt, a_0 - da, b_0, 35000)$ equals to

$$\left(\frac{7733069351623}{3125000000000}, \frac{25787091}{20000000000000}, \frac{189061375789453}{200000000000000}, -\frac{44841643}{200000000000000}, \frac{30543236182739}{250000000000000} \right)$$

and

$$\tilde{H}_W(t_0 - dt, 35000) < \frac{94851}{10000000000000}$$

Therefore,

$$\dot{F}(t_0 - dt, a_0 - da, b_0) > \frac{25787091}{200000000000000} - \frac{94851}{10000000000000} > 0$$

As pointed out above, we conclude that $\dot{F}(t_0 - dt, a, b_0) > 0$ for all $a \in [a_0 - da, a_0 + da]$. In the same way we have that $\dot{F}(t_0 + dt, a, b_0) < 0$ for all $a \in [a_0 - da, a_0 + da]$ because a direct computation shows that $Z_W(t_0 + dt, a_0 + da, b_0, 120000)$ equals to

$$\left(\frac{1546614123963}{6250000000000}, -\frac{198811}{6250000000000}, \frac{945307243792047}{100000000000000}, \frac{16995193}{20000000000000}, \frac{61086532113189}{50000000000000} \right)$$

and

$$\tilde{H}_W(t_0 + dt, 120000) < \frac{5651}{2000000000000}$$

Therefore,

$$\dot{F}(t_0 + dt, a_0 + da, b_0) < -\frac{198811}{62500000000000} + \frac{5651}{2000000000000} < 0$$

Let us show that $\dot{R}(t, a_0 - da, b_0) < 0$ for all $t \in [t_0 - dt, t_0 + dt]$. A direct computation shows that $Z_W(t_0 + dt, a_0 - da, b_0, 35000)$ equals to

$$\left(\frac{123729109625969}{500000000000000}, -\frac{196972647}{100000000000000}, \frac{945306878946787}{100000000000000}, -\frac{19027313}{250000000000000}, \frac{122173137972697}{1000000000000000} \right)$$

and

$$\tilde{H}_W(t_0 + dt, 35000) < \frac{94851}{100000000000000}$$

Therefore,

$$\dot{R}(t_0 + dt, a_0 - da, b_0) < -\frac{19027313}{250000000000000} + \frac{94851}{100000000000000} < 0$$

Since, $\ddot{R}(t, a_0 - da, b_0) > 0$ (see Corollary 7.10), we conclude that $\dot{R}(t, a_0 - da, b_0) < 0$ for all $t \in [t_0 - dt, t_0 + dt]$. Let us show that $\dot{R}(t, a_0 + da, b_0) > 0$ for all $t \in [t_0 - dt, t_0 + dt]$. A direct computation shows that $Z_W(t_0 - dt, a_0 + da, b_0, 35000)$ equals to

$$\left(\frac{24745827990179}{100000000000000}, \frac{55883369}{250000000000000}, \frac{189061484706171}{200000000000000}, \frac{52393253}{500000000000000}, \frac{61086475049353}{500000000000000} \right)$$

and

$$\tilde{H}_W(t_0 - dt, 35000) < \frac{94851}{100000000000000}$$

Therefore,

$$\dot{R}(t_0 - dt, a_0 + da, b_0) < \frac{52393253}{500000000000000} - \frac{94851}{100000000000000} > 0$$

Since, $\ddot{R}(t, a_0 + da, b_0) > 0$ (see Corollary 7.10), we conclude that $\dot{R}(t, a_0 + da, b_0) > 0$ for all $t \in [t_0 - dt, t_0 + dt]$. This finishes het proof of the Lemma. $\square$

Remark 7.12. *As a Corollary, using the Poincare-Miranda theorem, we obtain a proof of Lemma 6.3.*

Lemma 7.13. $\dot{F}(t_0 - st - dt, a, b_0 - sb) > 0$ for all $a \in [a_0 + sa - da, a_0 + sa + da]$, $\dot{F}(t_0 - st + dt, a, b_0 - sb) < 0$ for all $a \in [a_0 + sa - da, a_0 + sa + da]$, $\dot{R}(t, a_0 + sa - da, b_0 - sb) < 0$ for all $t \in [t_0 - st - dt, t_0 - st + dt]$ and $\dot{R}(t, a_0 + sa + da, b_0 - sb) > 0$ for all $t \in [t_0 - st - dt, t_0 - st + dt]$. Moreover, for all $t \in [t_0 - st - dt, t_0 - st + dt]$ and all $a \in [a_0 + sa - da, a_0 + sa + da]$, $\Theta(t, a, b_0 - sb) < \frac{7\pi}{18}$.

Proof. The proof is similar to that of Lemma 7.11. In this case $\dot{F}(t_0 - st - dt, a, b_0 - sb) > 0$ for all $a \in [a_0 + sa - da, a_0 + sa + da]$ because $Z_W(t_0 - st - dt, a_0 + sa - da, b_0 - sb, 35000)$ equals to

$$\left(\frac{247454580109467}{10000000000000}, \frac{14064311}{5000000000000}, \frac{945308716843341}{10000000000000}, -\frac{27791851}{5000000000000}, \frac{7635805749249}{6250000000000} \right)$$

and

$$\tilde{H}_W(t_0 - st - dt, 35000) < \frac{94849}{10000000000000}$$

Therefore,

$$\dot{F}(t_0 - st - dt, a_0 + sa - da, b_0 - sb) > \frac{14064311}{5000000000000} - \frac{94849}{10000000000000} > 0.$$

$\dot{F}(t_0 - st + dt, a, b_0 - sb) < 0$ for all $a \in [a_0 + sa - da, a_0 + sa + da]$, because $Z_W(t_0 - st + dt, a_0 + sa + da, b_0 - sb, 120000)$ equals to

$$\left(\frac{49490920135587}{20000000000000}, -\frac{4841111}{10000000000000}, \frac{29540903186787}{3125000000000}, \frac{21447973}{2500000000000}, \frac{122172932415207}{10000000000000} \right)$$

and

$$\tilde{H}_W(t_0 - st + dt, 120000) < \frac{5651}{2000000000000}$$

Therefore,

$$\dot{F}(t_0 - st + dt, a_0 + sa + da, b_0 - sb) < -\frac{4841111}{10000000000000} + \frac{5651}{2000000000000} < 0.$$

$\dot{R}(t, a_0 + sa + da, b_0 - sb) > 0$ for all $t \in [t_0 - st - dt, t_0 - st + dt]$, because $Z_W(t_0 - st - dt, a_0 + sa + da, b_0 - sb, 35000)$ equals to

$$\left(\frac{123727300365019}{50000000000000}, \frac{12058321}{20000000000000}, \frac{189061780400321}{20000000000000}, \frac{11254969}{20000000000000}, \frac{61086446906439}{50000000000000} \right)$$

and

$$\tilde{H}_W(t_0 - st - dt, 35000) < \frac{94849}{10000000000000}$$

Therefore,

$$\dot{R}(t_0 - st - dt, a_0 + sa + da, b_0 - sb) > \frac{11254969}{20000000000000} - \frac{94849}{10000000000000} > 0$$

$\dot{R}(t, a_0 + sa - da, b_0 - sb) < 0$ for all $t \in [t_0 - st - dt, t_0 - st + dt]$, because $Z_W(t_0 - st + dt, a_0 + sa - da, b_0 - sb, 35000)$ equals to

$$\left(\frac{4949091602187}{20000000000000}, -\frac{18526257}{50000000000000}, \frac{945308716843239}{100000000000000}, -\frac{25964629}{100000000000000}, \frac{122172930636361}{100000000000000} \right)$$

and

$$\tilde{H}_W(t_0 - st - dt, 35000) < \frac{1897}{200000000000}$$

Therefore,

$$\dot{R}(t_0 - st + dt, a_0 + sa - da, b_0 - sb) < -\frac{25964629}{100000000000000} + \frac{1897}{200000000000} < 0.$$

In order to show that $\Theta(t, a, b_0 - sb) < \frac{7\pi}{18}$, we first noticed that $Z_W(t_0 - st, a_0 + sa, b_0 - sb, 35000)$ equals to

$$\left(\frac{247454590419723}{100000000000000}, \frac{1161959}{100000000000000}, \frac{945308809422539}{100000000000000}, \frac{473597}{31250000000000}, \frac{122172912224601}{100000000000000} \right)$$

and

$$\tilde{H}_W(t_0 - st, 35000) < \frac{94849}{10000000000000}$$

Therefore,

$$\begin{aligned} \Theta(t_0 - st, a_0 + sa, b_0 - sb) - \frac{7\pi}{18} &< \frac{122172912224601}{100000000000000} + \frac{94849}{10000000000000} - \frac{7\pi}{18} \\ &= \frac{122172921709501}{100000000000000} - \frac{7\pi}{18} \end{aligned}$$

Using the mean value theorem, and the bounds $|\Theta_a| < \frac{483453}{1000000}$ and $|\dot{\Theta}| < \frac{27}{1000}$ found in Corollaries 7.10 and 7.6, we obtain that for any $t \in [t_0 - st - dt, t_0 - st + dt]$ and $a \in [a_0 + sa - da, a_0 + sa + da]$

$$\begin{aligned} \Theta(t, a, b_0 - sb) - \frac{7\pi}{18} &= \Theta(t, a, b_0 - sb) - \Theta(t, a_0 - sa, b_0 - sb) + \\ &\quad \Theta(t, a_0 + sa, b_0 - sb) - \Theta(t_0, a_0 + sa, b_0 - sb) + \\ &\quad \Theta(t_0, a_0 + sa, b_0 - sb) - \frac{7\pi}{18} \\ &< \frac{483453}{1000000} da + \frac{27}{1000} dt + \frac{122172921709501}{100000000000000} - \frac{7\pi}{18} \\ &< 0 \end{aligned}$$

□

Remark 7.14. As a Corollary, using the Poincare-Miranda theorem, we obtain a proof of Lemma 6.4

Lemma 7.15. $\dot{F}(t_0 + st - dt, a, b_0 + sb) > 0$ for all $a \in [a_0 - sa - da, a_0 - sa + da]$, $\dot{F}(t_0 + st + dt, a, b_0 + sb) < 0$ for all $a \in [a_0 - sa - da, a_0 - sa + da]$, $\dot{R}(t, a_0 - sa - da, b_0 + sb) < 0$ for all $t \in [t_0 + st - dt, t_0 + st + dt]$ and $\dot{R}(t, a_0 - sa + da, b_0 + sb) > 0$ for all $t \in [t_0 + st - dt, t_0 + st + dt]$. Moreover, for all all $t \in [t_0 + st - dt, t_0 + st + dt]$ and all $a \in [a_0 - sa - da, a_0 - sa + da]$, $\Theta(t, a, b_0 + sb) > \frac{7\pi}{18}$.

Proof. The proof is similar to that of Lemma 7.13. In this case $\dot{F}(t_0 + st - dt, a, b_0 + sb) > 0$ for all $a \in [a_0 - sa - da, a_0 - sa + da]$ because $Z_W(t_0 + st - dt, a_0 - sa - da, b_0 + sb, 35000)$ equals to

$$\left(\frac{247461898442221}{100000000000000}, \frac{1573331}{5000000000000}, \frac{189061080098229}{20000000000000}, -\frac{57188327}{100000000000000}, \frac{15271644451087}{125000000000000} \right)$$

and

$$\tilde{H}_W(t_0 + st - dt, 35000) < \frac{23713}{250000000000}$$

Therefore,

$$\dot{F}(t_0 - st - dt, a_0 + sa - da, b_0 - sb) > \frac{1573331}{5000000000000} - \frac{23713}{250000000000} > 0.$$

$\dot{F}(t_0 + st + dt, a, b_0 + b) < 0$ for all $a \in [a_0 + sa - da, a_0 + sa + da]$, because $Z_W(t_0 + st + dt, a_0 - sa + da, b_0 + sb, 120000, 15)$ equals to

$$\left(\frac{1237309595658901}{500000000000000}, -\frac{904301}{62500000000000}, \frac{9453055857540481}{1000000000000000}, \frac{842419313}{1000000000000000}, \frac{244346392152949}{200000000000000} \right)$$

and

$$\tilde{H}_W(t_0 + st + dt, 120000, 15) < \frac{77}{8000000000}$$

Therefore,

$$\dot{F}(t_0 - st + dt, a_0 + sa + da, b_0 - sb) < -\frac{904301}{62500000000000} + \frac{77}{8000000000} < 0.$$

$\dot{R}(t, a_0 - sa + da, b_0 + sb) > 0$ for all $t \in [t_0 + st - dt, t_0 + st + dt]$, because $Z_W(t_0 + st - dt, a_0 - sa + da, b_0 + sb, 35000)$ equals to

$$\left(\frac{49492383812687}{20000000000000}, \frac{63630339}{10000000000000}, \frac{945305585649929}{10000000000000}, \frac{1366741}{2500000000000}, \frac{24434631486741}{20000000000000} \right)$$

and

$$\tilde{H}_W(t_0 + st - dt, 35000) < \frac{23713}{250000000000}$$

Therefore,

$$\dot{R}(t_0 - st - dt, a_0 + sa + da, b_0 - sb) > \frac{1366741}{2500000000000} - \frac{23713}{250000000000} > 0$$

$\dot{R}(t, a_0 + sa - da, b_0 - sb) < 0$ for all $t \in [t_0 + st - dt, t_0 + st + dt]$, because $Z_W(t_0 + st + dt, a_0 - sa - da, b_0 + sb, 35000)$ equals to

$$\left(\frac{123730949221087}{50000000000000}, -\frac{33715619}{100000000000000}, \frac{945305400490959}{100000000000000}, -\frac{6891931}{25000000000000}, \frac{1908956160269}{1562500000000} \right)$$

and

$$\tilde{H}_W(t_0 + st + dt, 35000) < \frac{23713}{250000000000}$$

Therefore,

$$\dot{R}(t_0 + st + dt, a_0 + sa - da, b_0 - sb) < -\frac{6891931}{25000000000000} + \frac{23713}{250000000000} < 0.$$

In order to show that $\Theta(t, a, b_0 + sb) > \frac{7\pi}{18}$, we first noticed that $Z_W(t_0 + st, a_0 - sa, b_0 + sb, 35000)$ equals to

$$\left(\frac{123730954376411}{50000000000000}, \frac{3739331}{25000000000000}, \frac{945305493070449}{100000000000000}, \frac{1355097}{10000000000000}, \frac{24434635169083}{20000000000000} \right)$$

and

$$\tilde{H}_W(t_0 + st, 35000) < \frac{23713}{250000000000}.$$

Therefore,

$$\begin{aligned}
\Theta(t_0 - st, a_0 + sa, b_0 - sb) - \frac{7\pi}{18} &> \frac{24434635169083}{20000000000000} - \frac{23713}{2500000000000} - \frac{7\pi}{18} \\
&= \frac{24434637066123}{20000000000000} - \frac{7\pi}{18}
\end{aligned}$$

Using the mean value theorem, and the bounds $|\Theta_a| < \frac{483453}{1000000}$ and $|\dot{\Theta}| < \frac{27}{1000}$ found on corollaries 7.10 and 7.6, we obtain that for any $t \in [t_0 + st - dt, t_0 + st + dt]$ and $a \in [a_0 - sa - da, a_0 - sa + da]$

$$\begin{aligned}
\Theta(t, a, b_0 - sb) - \frac{7\pi}{18} &= \Theta(t, a, b_0 - sb) - \Theta(t, a_0 - sa, b_0 - sb) + \\
&\quad \Theta(t, a_0 + sa, b_0 - sb) - \Theta(t_0, a_0 + sa, b_0 - sb) + \\
&\quad \Theta(t_0, a_0 + sa, b_0 - sb) - \frac{7\pi}{18} \\
&> \frac{24434637066123}{20000000000000} - \frac{7\pi}{18} - \frac{483453}{1000000} da - \frac{27}{1000} dt \\
&> 0
\end{aligned}$$

□

Lemma 7.16. Let $\delta_1 = \frac{1017}{1250}$, $\tilde{\delta}_1 = \frac{8159}{10000}$, $\epsilon_1 = \frac{2677}{5000}$, $\delta_2 = \frac{16181}{10000}$, $\tilde{\delta}_2 = \frac{8359}{5000}$, $\epsilon_2 = \frac{4549}{5000}$, $\tilde{\epsilon}_1 = 6(st + dt)$, $\tilde{\epsilon}_2 = 3(sa + da)$, $\mu_1 = dt$ and $\mu_2 = da$. If $\rho_1 = \frac{\tilde{\epsilon}_1 - \mu_1}{m_1}$, $\rho_2 = \frac{\tilde{\epsilon}_2 - \mu_2}{m_2}$, where $m_1 = \frac{\epsilon_1(\tilde{\delta}_2 + \epsilon_2)}{\delta_1\delta_2 - \epsilon_1\epsilon_2}$ and $m_2 = \frac{\epsilon_2(\tilde{\delta}_1 + \epsilon_1)}{\delta_1\delta_2 - \epsilon_1\epsilon_2}$, then, for every $t \in [t_0 - \tilde{\epsilon}_1, t_0 + \tilde{\epsilon}_1]$, $a \in [a_0 - \tilde{\epsilon}_2, a_0 + \tilde{\epsilon}_2]$ and $b \in [b_0 - sb, b_0 + sb]$ we have that

$$\delta_1 < |\ddot{F}(at, a, b)| < \tilde{\delta}_1, \quad |\dot{F}_a(t, a, b)| < \epsilon_1, \quad |\dot{F}_b(t, a, b)| < \epsilon_1$$

and

$$|\ddot{R}(t, a, b)| < \epsilon_2, \quad \delta_2 < |\dot{R}_a(t, a, b)| < \tilde{\delta}_2, \quad |\dot{R}_b(t, a, b)| < \epsilon_2$$

Moreover, $sb < \rho_1$ and $sb < \rho_2$.

Proof. This is a direct computation. It uses Lemmas 7.5, 7.7 and 7.10

□

Remark 7.17. The previous lemma, along with Theorem 3.1 implies Lemma 6.6.

Remark 7.18. *Since we have shown the existence of a curve of initial conditions that provides reduced periodic functions, then we have not only shown the existence of the periodic solution with $\theta(t_0) = \frac{7\pi}{18}$ but we have shown the existence of infinitely many periodic solutions due to the fact that the function θ is not constant along this curve, and, on any open interval, there are infinitely many numbers of the form $\frac{p\pi}{q}$ with p and q whole numbers.*

8 Dictionary of functions

In this section we define the functions involved in the definition of the differential equations that we considered in this paper. The domain of the variable x_1 will be an interval containing possible values of the function $F(t, a, b)$. In the same way, the variable x_2 is related with the function $\dot{F}(t, a, b)$; the variable x_3 is related with the function $R(t, a, b)$; the variable x_4 is related with the function $\dot{R}(t, a, b)$; the variable x_5 is related with the function $F_a(t, a, b)$; the variable x_6 is related with the function $\dot{F}_a(t, a, b)$; the variable x_7 is related with the function $R_a(t, a, b)$; the variable x_8 is related with the function $\dot{R}_a(t, a, b)$; the variable x_9 is related with the function $F_b(t, a, b)$; the variable x_{10} is related with the function $\dot{F}_b(t, a, b)$; the variable x_{11} is related with the function $R_b(t, a, b)$; the variable x_{12} is related with the function $\dot{R}_b(t, a, b)$. As in the previous section, $s = \sqrt{4x_1^2 + x_3^2}$. For each one of these function $B(\phi_i) = \{\text{lb}_i, \text{ub}_i\}$ where lb_i is a lower bound and ub_i is a lower bound for the function $\phi_i = \phi_i(x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9, x_{10}, x_{11}, x_{12}, a)$ on the set V , where

$$V = I_1 \times I_2 \times I_3 \times I_4 \times I_5 \times I_6 \times I_7 \times I_8 \times I_9 \times I_{10} \times I_{11} \times I_{12} \times I_{13}$$

with

$$I_1 = \left[-\frac{1}{100}, \frac{62}{25}\right] \quad I_2 = \left[-\frac{1}{100}, \frac{3}{2}\right] \quad I_3 = \left[\frac{189}{20}, \frac{1001}{100}\right] \quad I_4 = \left[-\frac{33}{100}, \frac{1}{100}\right] \quad I_5 = \left[-\frac{1}{100}, \frac{31}{100}\right]$$

$$I_6 = \left[-\frac{1}{100}, \frac{12}{25}\right] \quad I_7 = \left[-\frac{1}{100}, \frac{69}{25}\right] \quad I_8 = \left[-\frac{1}{100}, \frac{42}{25}\right] \quad I_9 = \left[-\frac{1}{100}, \frac{101}{50}\right] \quad I_{10} = \left[\frac{1}{2}, \frac{101}{100}\right]$$

$$I_{11} = \left[-\frac{1}{100}, \frac{81}{100}\right] \quad I_{12} = \left[-\frac{1}{100}, \frac{89}{100}\right] \quad I_{13} = \left[\frac{43170106052787}{10000000000000}, \frac{43170844652787}{10000000000000}\right]$$

We would like to point out that $I_{13} = [a_0 - 3(sa + da), a_0 + 3(sa + da)]$.

Remark 8.1. *In order to obtain each bound $B(\phi_i) = \{lb_i, ub_i\}$, we first compute the minimum and maximum of the function, mi and ma respectively. Then we define $lb_i = \frac{1}{10^k} \lfloor 10^k mi \rfloor$ and $ub_i = \frac{1}{10^k} \lceil 10^k ma \rceil$, where k is a positive integer. Usually k is between 4 and 12 and it is chosen according to the precision that we want for the bound. The reason for using lb_i, ub_i and not the minimum and maximum is that ma and mi may have complicate expression in term of radicals and roots of polynomials.*

Remark 8.2. *The bounds for each of the functions for which we are writing an expression in terms of the variables $x_1, \dots, x_{12}$ have been obtained by directly using the Lagrange multipliers method. For those functions ϕ_i that we are written in term of the previous ϕ_j and some x_k , we used the method of Lagrange multiplier by changing each ϕ_j by a variable u_j and we used the fact that u_j has bound given by $B(\phi_j)$.*

$$8.1 \quad \phi_1 = -\frac{400x_1}{s^3}$$

Entry 2 of the vector field W . $B(\phi_1) = \left\{ -\frac{102003}{125000}, \frac{237}{50000} \right\}$

$$8.2 \quad \phi_2 = \frac{100a^2}{x_3^3} - \frac{25}{x_3^2} - \frac{200x_3}{s^3}$$

Entry 4 of the vector field W . $B(\phi_2) = \left\{ -\frac{387429}{1000000}, \frac{373771}{1000000} \right\}$

$$8.3 \quad \phi_3 = \frac{10a}{x_3^2}$$

Entry 5 of the vector field W . $B(\phi_3) = \left\{ \frac{215419}{500000}, \frac{483423}{1000000} \right\}$

$$8.4 \quad \phi_4 = -\frac{400(x_3^2 - 8x_1^2)}{s^5}$$

Entry (2, 1) of the matrix dW . $B(\phi_4) = \left\{ -\frac{94797}{200000}, -\frac{115837}{1000000} \right\}$

$$\mathbf{8.5} \quad \phi_5 = \frac{1200x_1x_3}{s^5}$$

Entry (2, 3) of the matrix dW . $B(\phi_5) = \left\{ -\frac{301}{200000}, \frac{101747}{500000} \right\}$

$$\mathbf{8.6} \quad \phi_6 = 50 \left(-\frac{6a^2}{x_3^4} + \frac{12x_3^2}{s^5} - \frac{4}{s^3} + \frac{1}{x_3^3} \right)$$

Entry (4, 3) of the matrix dW . $B(\phi_6) = \left\{ -\frac{83881}{200000}, -\frac{108213}{1000000} \right\}$

$$\mathbf{8.7} \quad \phi_7 = -\frac{20a}{x_3^3}$$

Entry (5, 3) of the matrix dW . $B(\phi_7) = \left\{ -\frac{12789}{125000}, -\frac{86081}{1000000} \right\}$

$$\mathbf{8.8} \quad \phi_8 = -\frac{1200(16x_1^2x_3 - x_3^3)}{s^7}$$

Part of the entry (2, 1) of the matrix dW_1 . $B(\phi_8) = \left\{ -\frac{3273}{500000}, \frac{18809}{125000} \right\}$

$$\mathbf{8.9} \quad \phi_9 = -\frac{4800(8x_1^3 - 3x_1x_3^2)}{s^7}$$

Part of the entry (2, 1) of the matrix dW_1 . $B(\phi_9) = \left\{ -\frac{1911}{1000000}, \frac{193837}{1000000} \right\}$

$$\mathbf{8.10} \quad \phi_{10} = x_4\phi_8 + x_2\phi_9$$

Entry (2, 1) of the matrix dW_1 . $B(\phi_{10}) = \left\{ -\frac{52523}{1000000}, \frac{73229}{250000} \right\}$

$$\mathbf{8.11} \quad \phi_{11} = \frac{4800(x_1^3 - x_1x_3^2)}{s^7}$$

Part of the entry (2, 3) of the matrix dW_1 . $B(\phi_{11}) = \left\{ -\frac{34393}{500000}, \frac{637}{1000000} \right\}$

$$\mathbf{8.12} \quad \phi_{12} = x_4\phi_{11} + x_2\phi_8$$

Entry (2, 3) of the matrix dW_1 . $B(\phi_{12}) = \left\{ -\frac{10507}{1000000}, \frac{31051}{125000} \right\}$

$$\mathbf{8.13} \quad \phi_{13} = \frac{1200a^2}{x_3^5} - \frac{3000x_3^3}{s^7} + \frac{1800x_3}{s^5} - \frac{150}{x_3^4}$$

Part of the Entry (4, 3) of the matrix dW_1 . $B(\phi_{13}) = \left\{ \frac{44031}{500000}, \frac{9611}{40000} \right\}$

$$\mathbf{8.14} \quad \phi_{14} = x_4\phi_{13} + 2x_2\phi_{11}$$

Entry (4, 3) of the matrix dW_1 . $B(\phi_{14}) = \left\{ -\frac{285649}{1000000}, \frac{2157}{500000} \right\}$

$$\mathbf{8.15} \quad \phi_{15} = \frac{60ax_4}{x_3^4}$$

Entry (5, 3) of the matrix dW_1 . $B(\phi_{15}) = \left\{ -\frac{10719}{1000000}, \frac{13}{40000} \right\}$

$$\mathbf{8.16} \quad \phi_{16} = \frac{400(8x_5x_1^2 + 3x_3x_7x_1 - x_3^2x_5)}{s^5}$$

Entry 6 of the vector field G . $B(\phi_{16}) = \left\{ -\frac{75543}{500000}, \frac{563009}{1000000} \right\}$

$$\mathbf{8.17} \quad \phi_{17} = -10\phi_7 + 2x_5\phi_5 + x_7\phi_6$$

Entry 8 of the vector field G . $B(\phi_{17}) = \left\{ -\frac{150409}{500000}, \frac{1153481}{1000000} \right\}$

$$\mathbf{8.18} \quad \phi_{18} = \frac{10(x_3 - 2ax_7)}{x_3^3}$$

Entry 9 of the vector field G . $B(\phi_{18}) = \left\{ -\frac{85201}{500000}, \frac{113003}{1000000} \right\}$

$$\mathbf{8.19} \quad \phi_{19} = x_5\phi_9 + x_7\phi_8$$

Entry (6, 1) of the matrix dG . $B(\phi_{19}) = \left\{ -\frac{10003}{500000}, \frac{475393}{1000000} \right\}$

$$\mathbf{8.20} \quad \phi_{20} = x_5\phi_8 + x_7\phi_{11}$$

Entry (6, 3) of the matrix dG . $B(\phi_{20}) = \left\{ -\frac{191879}{1000000}, \frac{9681}{200000} \right\}$

$$\mathbf{8.21} \quad \phi_{21} = -\frac{600a}{x_3^4}$$

Part of the entry (8, 3) of the matrix dG . $B(\phi_{21}) = \left\{ -\frac{324799}{1000000}, -\frac{257987}{1000000} \right\}$

$$\mathbf{8.22} \quad \phi_{22} = \phi_{21} + 2x_5\phi_1 + x_7\phi_{13}$$

Entry (8, 3) of the matrix dG . $B(\phi_{22}) = \left\{ -\frac{147363}{1000000}, \frac{289227}{500000} \right\}$

$$\mathbf{8.23} \quad \phi_{23} = \frac{60ax_7}{x_3^4} - \frac{20}{x_3^3}$$

Entry (9, 3) of the matrix dG . $B(\phi_{23}) = \left\{ -\frac{961}{40000}, \frac{32973}{500000} \right\}$

$$\mathbf{8.24} \quad \phi_{24} = x_8\phi_8 + x_6\phi_9$$

Part of the entry (6, 1) of the matrix dG_1 . $B(\phi_{24}) = \left\{ -\frac{1617}{125000}, \frac{69167}{200000} \right\}$

$$\mathbf{8.25} \quad \phi_{25} = -\frac{4800(16x_1^4 - 27x_3^2x_1^2 + x_3^4)}{s^9}$$

Part of the entry (6, 1) of the matrix dG_1 . $B(\phi_{25}) = \left\{ -\frac{15923}{250000}, \frac{8349}{500000} \right\}$

$$\mathbf{8.26} \quad \phi_{26} = \frac{24000(16x_1^3x_3 - 3x_1x_3^3)}{s^9}$$

Part of the entry (6, 1) of the matrix dG_1 . $B(\phi_{26}) = \left\{ -\frac{85577}{1000000}, \frac{1011}{1000000} \right\}$

$$\mathbf{8.27} \quad \phi_{27} = x_4\phi_{25} + x_2\phi_{26}$$

Part of the entry (6, 1) of the matrix dG_1 . $B(\phi_{27}) = \left\{ -\frac{33469}{250000}, \frac{4507}{200000} \right\}$

$$\mathbf{8.28} \quad \phi_{28} = \frac{4800(128x_1^4 - 96x_3^2x_1^2 + 3x_3^4)}{s^9}$$

Part of the entry (6, 1) of the matrix dG_1 . $B(\phi_{28}) = \left\{ -\frac{64021}{1000000}, \frac{7643}{40000} \right\}$

$$\mathbf{8.29} \quad \phi_{29} = x_2\phi_{28} + x_4\phi_{26}$$

Part of the entry (6, 1) of the matrix dG_1 . $B(\phi_{29}) = \left\{ -\frac{12111}{125000}, \frac{314853}{1000000} \right\}$

$$\mathbf{8.30} \quad \phi_{30} = \phi_{24} + x_7\phi_{27} + x_5\phi_{29}$$

Entry (6, 1) of the matrix dG_1 . $B(\phi_{30}) = \left\{ -\frac{41247}{100000}, \frac{505637}{1000000} \right\}$

$$\mathbf{8.31} \quad \phi_{31} = x_6\phi_8 + x_8\phi_{11}$$

Part of the entry (6, 3) of the matrix dG_1 . $B(\phi_{31}) = \left\{ -\frac{118703}{1000000}, \frac{73297}{1000000} \right\}$

$$\mathbf{8.32} \quad \phi_{32} = -\frac{24000(3x_1^3x_3 - x_1x_3^3)}{s^9}$$

Part of the entry (6, 3) of the matrix dG_1 . $B(\phi_{32}) = \left\{ -\frac{337}{1000000}, \frac{6067}{200000} \right\}$.

$$\mathbf{8.33} \quad \phi_{33} = x_2\phi_{25} + x_4\phi_{32}$$

Part of the entry (6, 3) of the matrix dG_1 . $B(\phi_{33}) = \left\{ -\frac{105549}{1000000}, \frac{25351}{1000000} \right\}$

$$\mathbf{8.34} \quad \phi_{34} = \phi_{31} + x_7\phi_{33} + x_5\phi_{27}$$

Entry (6, 3) of the matrix dG_1 . $B(\phi_{34}) = \left\{ -\frac{1411}{3125}, \frac{37563}{250000} \right\}$

$$\mathbf{8.35} \quad \phi_{35} = \frac{2400ax_4}{x_3^5}$$

Part of the entry (8, 3) of the matrix dG_1 . $B(\phi_{35}) = \left\{ -\frac{45369}{1000000}, \frac{11}{8000} \right\}$

$$\mathbf{8.36} \quad \phi_{36} = \phi_{35} + x_8\phi_{13} + 2x_6\phi_{11}$$

Part of the entry $(8, 3)$ of dG_1 . $B(\phi_{36}) = \left\{ -\frac{113807}{1000000}, \frac{406413}{1000000} \right\}$

$$\mathbf{8.37} \quad \phi_{37} = 600 \left(-\frac{10a^2}{x_3^6} + \frac{8x_3^4}{s^9} - \frac{108x_1^2x_3^2}{s^9} + \frac{12x_1^2}{s^7} + \frac{1}{x_3^5} \right)$$

Part of the entry $(8, 3)$ of the matrix dG_1 . $B(\phi_{37}) = \left\{ -\frac{579}{4000}, -\frac{2871}{50000} \right\}$

$$\mathbf{8.38} \quad \phi_{38} = x_4\phi_{37} + 2x_2\phi_{32}$$

Part of the entry $(8, 3)$ of the matrix dG_1 . $B(\phi_{38}) = \left\{ -\frac{2459}{1000000}, \frac{138773}{1000000} \right\}$

$$\mathbf{8.39} \quad \phi_{39} = \phi_{36} + 2x_5\phi_{33} + x_7\phi_{38}$$

Entry $(8, 3)$ of the matrix dG_1 . $B(\phi_{39}) = \left\{ -\frac{117597}{1000000}, \frac{400107}{1000000} \right\}$

$$\mathbf{8.40} \quad \phi_{40} = \frac{60(x_3(ax_8 + x_4) - 4ax_4x_7)}{x_3^5}$$

Entry $(9, 3)$ of the matrix dG_1 . $B(\phi_{40}) = \left\{ -\frac{2853}{1000000}, \frac{32303}{500000} \right\}$

$$\mathbf{8.41} \quad \phi_{41} = \frac{400(8x_9x_1^2 + 3x_3x_{11}x_1 - x_3^2x_9)}{s^5}$$

Entry 6 of the vector field U . $B(\phi_{41}) = \left\{ -\frac{19173}{20000}, \frac{83109}{500000} \right\}$

$$\mathbf{8.42} \quad \phi_{42} = 2x_9\phi_5 + x_{11}\phi_6$$

Entry 8 of the vector field U . $B(\phi_{42}) = \left\{ -\frac{19549}{500000}, \frac{205701}{250000} \right\}$

$$\mathbf{8.43} \quad \phi_{43} = x_{11}\phi_8 + x_9\phi_9$$

Entry $(6, 1)$ of the matrix dU . $B(\phi_{43}) = \left\{ -\frac{9163}{1000000}, \frac{256717}{500000} \right\}$

$$\mathbf{8.44} \quad \phi_{44} = x_{11}\phi_{11} + x_9\phi_8$$

Entry (6, 3) of the matrix dU . $B(\phi_{44}) = \left\{ -\frac{3447}{50000}, \frac{152321}{500000} \right\}$

$$\mathbf{8.45} \quad \phi_{45} = 2x_9\phi_{11} + x_{11}\phi_{13}$$

Entry (8, 3) of the matrix dU . $B(\phi_{45}) = \left\{ -\frac{280299}{1000000}, \frac{197197}{1000000} \right\}$

$$\mathbf{8.46} \quad \phi_{46} = x_{12}\phi_8 + x_{10}\phi_9$$

Part of the entry (6, 1) of the vector field dU_1 . $B(\phi_{46}) = \left\{ -\frac{7757}{1000000}, \frac{10303}{31250} \right\}$

$$\mathbf{8.47} \quad \phi_{47} = x_4\phi_{26} + x_2\phi_{28}$$

Part of the entry (6, 1) of the matrix dU_1 . $B(\phi_{47}) = \left\{ -\frac{12111}{125000}, \frac{314853}{1000000} \right\}$

$$\mathbf{8.48} \quad \phi_{48} = x_{11}\phi_{27} + x_9\phi_{47} + \phi_{46}$$

Entry (6, 1) of the matrix dU_1 . $B(\phi_{48}) = \left\{ -\frac{203233}{500000}, \frac{35511}{62500} \right\}$

$$\mathbf{8.49} \quad \phi_{49} = x_{12}\phi_{11} + x_{10}\phi_8$$

Part of the entry (6, 3) of the matrix dU_1 . $B(\phi_{49}) = \left\{ -\frac{67831}{1000000}, \frac{30533}{200000} \right\}$

$$\mathbf{8.50} \quad \phi_{50} = x_9\phi_{27} + x_{11}\phi_{33} + \phi_{49}$$

Entry (6, 3) of the matrix dU_1 . $B(\phi_{50}) = \left\{ -\frac{105939}{250000}, \frac{218721}{1000000} \right\}$

$$\mathbf{8.51} \quad \phi_{51} = \frac{1200a^2}{x_3^5} - \frac{1200x_3^5}{s^9} + \frac{2400x_1^2x_3^3}{s^9} + \frac{28800x_1^4x_3}{s^9} - \frac{150}{x_3^4}$$

Part of the entry (8, 3) of the matrix dU_1 . $B(\phi_{51}) = \left\{ \frac{44031}{500000}, \frac{9611}{40000} \right\}$

$$\mathbf{8.52} \quad \phi_{52} = 2x_{10}\phi_{11} + x_{12}\phi_{51}$$

Part of the entry (8, 3) of the matrix dU_1 . $B(\phi_{52}) = \left\{ -\frac{141351}{1000000}, \frac{53783}{250000} \right\}$

$$\mathbf{8.53} \quad \phi_{53} = 2x_9\phi_{33} + x_{11}\phi_{38} + \phi_{52}$$

Entry (8, 3) of the matrix dU_1 . $B(\phi_{53}) = \left\{ -\frac{569761}{1000000}, \frac{429957}{1000000} \right\}$

$$\mathbf{8.54} \quad \phi_{54} = \frac{100(a - a_0)(a + a_0)}{x_3^3}$$

Entry 4 of $W - W_0$. $B(\phi_{54}) = \left\{ -\frac{19}{500000}, \frac{19}{500000} \right\}$

$$\mathbf{8.55} \quad \phi_{55} = \frac{10(a - a_0)}{x_3^2}$$

Entry 5 of $W - W_0$. $B(\phi_{55}) = \left\{ -\frac{1}{200000}, \frac{1}{200000} \right\}$

$$\mathbf{8.56} \quad \phi_{56} = -\frac{100(a - a_0)(3ax_7 + 3a_0x_7 - 2x_3)}{x_3^4}$$

Entry 8 of $G - G_0$. $B(\phi_{56}) = \left\{ -\frac{1}{40000}, \frac{1}{40000} \right\}$

$$\mathbf{8.57} \quad \phi_{57} = -\frac{20x_7(a - a_0)}{x_3^3}$$

Entry 9 of $G - G_0$. $B(\phi_{57}) = \left\{ -\frac{3}{1000000}, \frac{3}{1000000} \right\}$

$$\mathbf{8.58} \quad \phi_{58} = -\frac{300x_{11}(a - a_0)(a + a_0)}{x_3^4}$$

Entry 8 of $U - U_0$. $B(\phi_{57}) = \left\{ -\frac{1}{1000000}, \frac{1}{1000000} \right\}$

References

- [1] Chenciner, A., Montgomery, R. *A remarkable periodic solution of the three body problem in the case of equal masses*. Annals of Math. **152** (2000), 881-901.
- [2] Gronwall, T. H. *Note on the Derivatives with respect to a parameter of Solutions of a System of Differential Equations*. Annals of Math. **20** (1919), 292-296.

- [3] Meyer, K. & Schmidt, D *Libration of central configurations and braided saturn rings*. Celestial Mechanics and Dynamical Astronomy **55**: 289-303, 1993.
- [4] C. Moore *Braids in Classical Gravity*. Physical Review Letters **70** 3675-3679.
- [5] H. Hénon *A family of periodic solutions of the planar three-body problem, and their stability*. Celestial Mechanics **13** (1976) pp. 267-285
- [6] Perdomo *A family of solution of the n body problem*. <http://arxiv.org/pdf/1410.1757.pdf>
- [7] Yan, D., Ouyang, T. *New phenomena in the spatial isosceles three-body problem*. <http://arxiv.org/abs/1404.4459>
- [8] C. Simo *Dynamical properties of the figure eight solution of the three-body problem*. Celestial mechanics (Evanston, IL, 1999), 209228, Contemp. Math., 292, Amer. Math. Soc., Providence, RI, 2002.